

DX0589K1B

PATENT APPLICATION

5

MAMMALIAN CHEMOKINE REAGENTS

INVENTORS:

10

Wei Wang, a citizen of the Peoples' Republic of China,  
residing at 3090 Middlefield Road; Palo Alto,  
California 94306;

15

Kurt C. Gish, a citizen of the United States, residing  
at 974 Hayes Street; San Francisco, California  
94117;

20

Thomas J. Schall, a citizen of the United States,  
residing at 2050 Mills Avenue; Menlo Park,  
California 94025;

25

Alain P. Vicari, a citizen of France, residing at 2250  
Latham Street #54; Mountain View, California  
94040; and

30

Albert Zlotnik, a citizen of the United States,  
residing at 507 Alger Drive; Palo Alto,  
California 94306.

EM161836205US

35

Express Mail label number EM 161 836 205 US  
Date of Deposit is July 3, 1997  
I hereby certify that this paper is being deposited with the United  
States Postal Service "Express Mail Post Office to Addressee" service  
under 37 C.F.R. § 1.10 on the date indicated above and is addressed to:  
Box Patent Application; Honorable Commissioner for Patents and  
Trademarks; Washington, D.C. 20231.

40

Lois E. Miller July 3, 1997  
Lois E. Miller date



28008

PATENT TRADEMARK OFFICE

45

Assignee:

Schering Corporation, a New Jersey corporation

EL789848704US

50

DNAX Research Institute  
901 California Avenue  
Palo Alto, California 94304-1104  
Tel: (415)852-9196  
Fax: (415)496-1200

"Express Mail" mailing label number  
Date of Deposit January 3, 2002  
I hereby certify that this paper or fee is being  
deposited with the United States Postal Service  
"Express Mail Post Office to Addressee" service  
under 37 C.F.R. § 1.10 on the date indicated above  
and is addressed to the Commissioner of Patents  
and Trademarks, Washington, D.C. 20231.

Lois E. Miller  
(Typed or printed name of person mailing) (Signature of person mailing) (Date)

## MAMMALIAN CHEMOKINE REAGENTS

This filing is a regular U.S. Patent Application resulting from the conversion of provisional applications.

5

## FIELD OF THE INVENTION

The present invention relates to compositions related to proteins which function in controlling physiology, development, and/or differentiation of mammalian cells, e.g., cells of a mammalian immune system. In particular, it provides proteins and mimetics which regulate physiology, development, differentiation, and function of various cell types, including hematopoietic cells. It also provides receptor reagents for chemokine-like proteins.

15

## BACKGROUND OF THE INVENTION

The circulating component of the mammalian circulatory system comprises various cell types, including red and white blood cells of the erythroid or the myeloid cell lineages. See, e.g., Rapaport (1987) Introduction to Hematology (2d ed.) Lippincott, Philadelphia, PA; Jandl (1987) Blood: Textbook of Hematology, Little, Brown and Co., Boston, MA.; and Paul (ed.) (1993) Fundamental Immunology 3d ed, Raven Press, N.Y. Progression through various stages of differentiation are regulated by various signals provided to the cells, often mediated through a class of proteins known as the cytokines. Within this group of molecules as a further group known as the chemoattractant cytokines, or chemokines. See, e.g., Schall (1994) "The Chemokines" in Thomson (ed.) The Cytokine Handbook (2d ed.) Academic Press; and Schall and Bacon (1994) Current Opinion in Immunology 6:865-873.

Although the full spectrum of biological activities of the chemokines has not been extensively investigated, chemoattractant effects are recognized. The best known biological functions of these molecules relate to chemoattraction of leukocytes. However, new chemokines are being discovered, and their biological effects on the

various cells responsible for immunological responses are topics of continued study.

Certain G-protein coupled receptors have also been characterized, presumably chemokine receptors. See, e.g.,  
5 Samson, et al. (1996) Biochemistry 35:3362-3367; and  
Rapport, et al. (1996) J. Leukocyte Biology 59:18-23.

These observations indicate that other factors exist whose functions in hematopoiesis, immune development, and leukocyte trafficking were heretofore unrecognized. These  
10 factors provide for biological activities whose spectra of effects are distinct from known differentiation, activation, or other signaling factors. The absence of knowledge about the structural, biological, and  
physiological properties of the regulatory factors which  
15 regulate hematopoietic cell physiology in vivo prevents the modification of the effects of such factors. Thus, medical conditions where regulation of the development or physiology of relevant cells is required remains unmanageable.

## SUMMARY OF THE INVENTION

The present invention is based, in part, upon the discovery of new genes encoding chemokines, and new genes encoding various receptors for chemokines. It embraces  
5 agonists and antagonists of the chemokines. In particular, sequences of various chemokines, e.g., designated Thymus Expressed ChemoKine (TECK); MIP-3 $\alpha$ ; MIP-3 $\beta$ ; and 7 transmembrane receptors, designated "dendritic cell receptor for chemokine" (DC CR) and "monocyte/dendritic  
10 cell receptor for chemokine" (M/DC CR); and mutations (muteins) of the respective natural sequences, fusion proteins, chemical mimetics, antibodies, and other structural or functional analogs are provided. It is also directed to isolated genes encoding respective proteins of  
15 the invention. Various uses of these different protein or nucleic acid compositions are also provided.

The present invention provides a substantially pure or isolated polypeptide comprising a segment exhibiting sequence homology to a corresponding portion of a mature  
20 TECK, MIP-3 $\alpha$ , MIP-3 $\beta$ , DC CR, or M/DC CR, wherein the homology is at least about 70% identity and the portion is at least about 25 amino acids. Preferably, the protein further comprises a second segment exhibiting at least about 90% identity over at least 9 amino acids; or at least  
25 about 80% identity over at least 17 amino acids. In other preferred embodiments, the polypeptide: is from a warm blooded animal selected from the group of birds and mammals, including a mouse or human; comprises a natural sequence from Tables 1 through 5; exhibits a post-  
30 translational modification pattern distinct from a natural form of the polypeptide; is made by expression of a recombinant nucleic acid; comprises synthetic sequence; is detectably labeled; is conjugated to a solid substrate; is conjugated to another chemical moiety; is a fusion protein;  
35 is in a denatured conformation, including detergent denaturation; further comprises an epitope tag; is an immunogenic polypeptide; has a defined homogeneous



molecular weight; is useful as a carbon source; is an allelic variant of SEQ ID NO: 2, 4, 6, 8, 10, or 12; is a 3-fold or less substituted form of a natural sequence; is in a sterile composition; is in a buffered solution or  
5 suspension; is in a regulated release device; comprises a post-translational modification; is in a cell; or is in a kit which further comprises instructions for use or disposal of reagents therein.

In other aspects, the invention provides an isolated  
10 or recombinant nucleic acid encoding such protein, where the portion consists of sequence from the coding region of SEQ ID NO: 1, 3, 5, 7, 9, or 11. Other aspects include such nucleic acids which: exhibit at least about 80% identity to a natural cDNA encoding said segment; is in an  
15 expression vector; further comprises a promoter; further comprises an origin of replication; is from a natural source; is detectably labeled; comprises synthetic nucleotide sequence; is less than 6 kb; is from a mammal; comprises a natural full length mature coding sequence; is  
20 in a kit, which also comprises instructions for use or disposal of reagents therein; is a specific hybridization probe for a gene encoding the protein; is a PCR product; or is in a cell. The invention also provides a method of using a purified nucleic acid by expressing the nucleic  
25 acid to produce a protein.

Alternatively, the invention provides an isolated or recombinant nucleic acid which encodes at least eight consecutive residues of SEQ ID NO: 2, 4, 6, 8, 10, or 12. Preferably, that nucleic acid encodes at least: twelve  
30 consecutive residues from SEQ ID NO: 2, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 1; twelve consecutive residues from SEQ ID NO: 4, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 3; twelve consecutive residues  
35 from SEQ ID NO: 6, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 5; twelve consecutive residues from SEQ ID NO: 8, and further

comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 7; twelve consecutive residues from SEQ ID NO: 10, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 9; or twelve consecutive  
5 residues from SEQ ID NO: 12, and further comprises a coding sequence of at least 17 nucleotides from SEQ ID NO: 11. In other preferred embodiments, the nucleic acid: exhibits at least about 80% identity to a natural cDNA encoding the segment; is in an expression vector; further comprises a  
10 promoter; further comprises an origin of replication; encodes a 3-fold or less substituted sequence from a natural sequence; is from a natural source; is detectably labeled; comprises synthetic nucleotide sequence; is less than 6 kb; is from a mammal; is attached to a solid  
15 substrate, including in a Southern or Northern blot; comprises a natural full length coding sequence; is in a cell; or is in a detection kit, which also comprises instructions for use or disposal of reagents therein. Further embodiments include a nucleic acid which hybridizes  
20 under stringent wash conditions of 55° C and less than 150 mM salt to the nucleic acid; while preferred embodiments include those which exhibit at least about 85% identity over a stretch of at least about 30 nucleotides to a primate sequence of SEQ ID NO: 1, 3, 5, 7, 9, or 11; or  
25 where the identity is at least 90%; or the stretch is at least 75 nucleotides; or where the identity is at least 95%; or the stretch is at least 100 nucleotides.

In other embodiments, the invention provides a binding compound comprising an antigen binding fragment from an  
30 antibody which binds to a mature TECK, MIP-3 $\alpha$ , MIP-3 $\beta$ , DC CR, or M/DC CR protein. In various embodiments, the binding compound is one wherein: the polypeptide is a mouse or human protein; the antibody is raised against a mature peptide sequence of Tables 1 through 5; the antibody is a  
35 monoclonal antibody; the binding compound is attached to a solid substrate; the binding compound is in a sterile composition; the binding compound binds to a denatured

antigen, including a detergent denatured antigen; the binding compound is detectably labeled; the binding compound is an Fv, Fab, or Fab2 fragment; the binding compound is conjugated to a chemical moiety; the binding  
5 compound is in a detection kit which also comprises instructions for use or disposal of reagents therein.

The invention also provides a cell which makes the antibody.

The invention embraces methods of purifying a  
10 polypeptide using a binding compound to specifically separate said polypeptides from others; of generating an antigen-binding compound complex comprising the step of contacting a sample comprising the antigen to a sample comprising a binding compound; or of modulating physiology  
15 or development of a cell expressing a receptor for a chemokine selected from TECK, MIP-3 $\alpha$ , or MIP-3 $\beta$ ; the method comprising contacting the cell with a composition comprising an agonist or mutein of said chemokine or an antibody antagonist of the chemokine. In certain  
20 embodiments of the method, the cell is a macrophage, lymphocyte, or eosinophil; or the physiology is a cellular calcium flux, a chemoattractant response, cellular morphology modification responses, phosphoinositide lipid turnover, or an antiviral response. In other embodiments,  
25 the receptor is DC CR and the chemokine is MIP-3 $\alpha$ , the physiology is pulmonary physiology, or the cell is an eosinophil.

## DESCRIPTION OF THE DRAWINGS

Figures 1A-1B show chemotactic properties of mTECK recombinant protein. Fig. 1A shows migration of mouse thymocytes to recombinant mTECK and effect of pertussis toxin. Chemotaxis assays were performed as described. Recombinant mouse lymphotactin was used as a positive control. Data are expressed as the mean of cell counts obtained from three separate experiments in duplicate  $\pm$  SEM. In one experiment, cells were pre incubated 1 h with 10 ng/ml pertussis toxin (PTX) prior to the assay. Fig. 1B shows migration of other leukocyte subsets to recombinant mTECK. Mouse splenic dendritic cells and mouse activated macrophages were obtained. THP-1 human monocytic cells were used without or with a 16 h activation with IFN- $\gamma$ . Results are obtained as the mean of the chemotactic index from three separate experiments per cell type in duplicate  $\pm$  SD. The number of cells migrating to medium alone was greater than 40 cells per 5 high power fields in each experiment. Recombinant MIP-1 $\alpha$  was used as a positive control.

## DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

## OUTLINE

- 5
- I. General
  - II. Purified Chemokines, Receptors
    - A. physical properties
    - B. biological properties
  - 10 III. Physical Variants
    - A. sequence variants, fragments
    - B. post-translational variants
      - 1. glycosylation
      - 2. others
  - 15 IV. Functional Variants
    - A. analogs; fragments
      - 1. agonists
      - 2. antagonists
    - B. mimetics
      - 20 1. protein
      - 2. chemicals
    - C. species variants
  - V. Antibodies
    - 25 A. polyclonal
    - B. monoclonal
    - C. fragments, binding compositions
  - VI. Nucleic Acids
    - 30 A. natural isolates; methods
    - B. synthetic genes
    - C. methods to isolate
  - VII. Making Chemokines, Receptors; Mimetics
    - A. recombinant methods
    - B. synthetic methods
    - C. natural purification
  - 35 VIII. Uses
    - A. diagnostic
    - B. therapeutic
  - IX. Kits
    - 40 A. nucleic acid reagents
    - B. protein reagents
    - C. antibody reagents
  - X. Receptors

## I. General

- 45 The present invention provides DNA sequences encoding various mammalian proteins which exhibit structural properties characteristic of a chemotactic cytokine, or chemokine. Other embodiments are directed to chemokine receptors. See, e.g., Lodi, et al. (1994) Science
- 50 263:1762-1767; Gronenborn and Clore (1991) Protein

- Engineering 4:263-269; Miller and Kranger (1992) Proc. Nat'l Acad. Sci. USA 89:2950-2954; Matsushima and Oppenheim (1989) Cytokine 1:2-13; Stoeckle and Baker (1990) New Biol. 2:313-323; Oppenheim, et al. (1991) Ann. Rev. Immunol. 9:617-648; Schall (1991) Cytokine 3:165-183; and The Cytokine Handbook Academic Press, NY. Mouse and human embodiments are described herein.

Chemokines play an important role in immune and inflammatory responses by inducing migration and adhesion of leukocytes. These small secreted molecules are a growing superfamily of 8-14 kDa proteins characterized by a conserved four cysteine motif. See, e.g., Schall (1991) Cytokine 3:165-183; and Thomson (ed.) The Cytokine Handbook Academic Press, NY. Chemokines are secreted by activated leukocytes and act as a chemoattractant for a variety of cells which are involved in inflammation. Besides chemoattractant properties, chemokines have been shown to induce other biological responses, e.g., modulation of second messenger levels such as  $Ca^{++}$ ; inositol phosphate pool changes (see, e.g., Berridge (1993) Nature 361:315-325 or Billah and Anthes (1990) Biochem. J. 269:281-291); cellular morphology modification responses; phosphoinositide lipid turnover; possible antiviral responses; and others. Thus, the chemokines provided herein may, alone or in combination with other therapeutic reagents, have advantageous combination effects.

Moreover, there are reasons to suggest that chemokines may have effects on other cell types, e.g., attraction or activation of monocytes, dendritic cells, T cells, eosinophils, and/or perhaps on basophils and/or neutrophils. They may also have chemoattractive effects on various neural cells including, e.g., dorsal root ganglia neurons in the peripheral nervous system and/or central nervous system neurons.

Membrane proteins which contain seven transmembrane segments have been characterized as G-protein coupled receptors. Many of these receptors have been characterized

as receptors for chemokines, based in part on structural features. Chemokine receptors are important in the signal transduction mechanisms mediated by the chemokines. They are useful markers for distinguishing cell populations, and have been implicated as specific receptors for retroviral infections.

The chemokine superfamily was classically divided into two groups exhibiting characteristic structural motifs, the Cys-X-Cys (C-X-C) and Cys-Cys (C-C) families. These were distinguished on the basis of a single amino acid insertion between the NH-proximal pair of cysteine residues and sequence similarity. Typically, the C-X-C chemokines, i.e., IL-8 and MGSA/Gro- $\alpha$  act on neutrophils but not on monocytes, whereas the C-C chemokines, i.e., MIP-1 $\alpha$  and RANTES, are potent chemoattractants for monocytes and lymphocytes but not neutrophils. See, e.g., Miller, et al. (1992) Crit. Rev. Immunol. 12:17-46. A recently isolated chemokine, lymphotactin, does not belong to either group and may constitute a first member of a third chemokine family, the C family. Lymphotactin does not have a characteristic CC or CXC motif, and acts on lymphocytes but not neutrophils and monocytes. See, e.g., Kelner et al. (1994) Science 266:1395-1399. This chemokine defines a new C-C chemokine family. Even more recently, another chemokine exhibiting a CX3C motif has been identified, which establishes a fourth structural class.

The present invention provides additional chemokine reagents, e.g., nucleic acids, proteins and peptides, antibodies, etc., related to the newly discovered respective chemokines designated TECK; MIP-3 $\alpha$ , and MIP-3 $\beta$ .

In other embodiments, the invention provides two genes encoding novel 7-transmembrane (7-TM) receptors, presumably G-protein coupled receptors and likely chemokine receptors. These 7-TM receptors are hypothesized to be chemokine receptors and have been designated DC CR and M/DC CR. Their ligands have not yet specifically been completely identified. However, the receptors exhibit structural

features typical of known chemokine receptors, e.g., 7  
transmembrane spanning structures. They may exhibit  
properties of binding many different cytokines at varying  
specificities (shared or promiscuous binding specificity)  
5 or may exhibit high affinity for one (specific) or a subset  
(shared) of chemokines.

The described chemokines and receptors should be  
important for mediating various aspects of cellular, organ,  
tissue, or organismal physiology or development.

## 10 II. Purified chemokines, receptors

Mouse and human Thymus Expressed ChemoKine (TECK)  
nucleotide and amino acid sequences are shown in Table 1.  
Nucleotide and amino acid sequences of another novel  
15 chemokine, from human, designated MIP-3 $\alpha$  are provided in  
Table 2. Nucleotide and derived amino acid sequences of a  
third novel chemokine, from human, designated MIP-3 $\beta$  are  
shown in Table 3. Generic descriptions of physical  
properties of polypeptides, nucleic acids, and antibodies  
20 where directed to one embodiment clearly are generally  
applicable to other chemokines or receptors described  
herein.

The nucleotide and amino acid sequences of a novel  
chemokine receptor found on dendritic cells (DC), from  
25 human, and designated DC CR, are provided in Table 4. The  
nucleotide and amino acid sequences of another novel  
chemokine receptor found on macrophages and dendritic  
cells, from human, and designated M/DC CR, are provided in  
Table 5.

30 These amino acid sequences, provided amino to carboxy,  
are important in providing sequence information on the  
chemokine ligand or receptor, allowing for distinguishing  
the protein from other proteins. Moreover, the sequences  
allow preparation of peptides to generate antibodies to  
35 recognize and distinguish such segments, and allow  
preparation of oligonucleotide probes, both of which are  
strategies for isolation, e.g., cloning, of genes encoding



such sequences, or related sequences, e.g., natural polymorphic or other variants. Similarities of the chemokines have been observed with other cytokines. See, e.g., Bosenberg, et al. (1992) Cell 71:1157-1165; Huang, et. al. (1992) Molecular Biology of the Cell 3:349-362; and  
5 Pandiella, et al. (1992) J. Biol. Chem. 267:24028-24033. Likewise for the receptors.

Table 1: Nucleotide sequence (5' to 3') of TECK from mouse and the corresponding amino acid sequence (amino to carboxy). Signal sequence probably runs as shown between Ala and Gln, see SEQ ID NO: 1 and 2. Human sequences are SEQ ID NO: 3 and 4.

5	1	AGGCTACAAGCAGGCACCAGCTCTCAGGACCAGAAAGGCATTGGTGGCCCCCTTAAACCT	60
10	61	TCAGGTATCTGGAGAGGAGATCTAACCTTCACTATGAAACTGTGGCTTTTTGCCTGCCTG	120
	1	MetLysLeuTrpLeuPheAlaCysLeu	9
	121	GTTGCCTGTTTTGTTGGGGCCTGGATGCCGGTTGTCCATGCCCAAGGTGCCTTTGAAGAC	180
	10	ValAlaCysPheValGlyAlaTrpMetProValValHisAlaGlnGlyAlaPheGluAsp	29
15	181	TGCTGCCTGGGTTACCAGCACAGGATCAAATGGAATGTTCTCCGGCATGCTAGGAATTAT	240
	30	CysCysLeuGlyTyrGlnHisArgIleLysTrpAsnValLeuArgHisAlaArgAsnTyr	49
	241	CACCAGCAGGAAGTGAGTGGAAGCTGCAACCTACGTGCTGTGAGATTCTACTTCCGCCAG	300
20	50	HisGlnGlnGluValSerGlySerCysAsnLeuArgAlaValArgPheTyrPheArgGln	69
	301	AAAGTAGTGTTGGGAATCCAGAGGACATGAATGTGAAGAGGGCGATAAGAATCTTGACA	360
	70	LysValValCysGlyAsnProGluAspMetAsnValLysArgAlaIleArgIleLeuThr	89
	361	GCTAGGAAAAGGCTAGTCCACTGGAAGAGCGCCTCAGACTCTCAGACTGAAAGGAAGAAG	420
25	90	AlaArgLysArgLeuValHisTrpLysSerAlaSerAspSerGlnThrGluArgLysLys	109
	421	TCAAACCATATGAAGTCCAAGGTGGAGAACCCCAACAGTACAAGCGTGAGGAGTGCCACC	480
	110	SerAsnHisMetLysSerLysValGluAsnProAsnSerThrSerValArgSerAlaThr	129
30	481	CTAGGTCATCCCAGGATGGTGATGATGCCCAGAAAGACCAACAATTAAGTTAATTACTCA	540
	130	LeuGlyHisProArgMetValMetMetProArgLysThrAsnAsnEnd	144
	541	GAGTAAGCACCAGCTGGAGGATGGGCGGAGTCTGCTGAAGTGCTGTCTTCTAGGCATGCC	600
35	601	AGTGCCAATGAACTCACTGAAGCTACAGTTTCCTGTACAAGACCAGACCCACCAACGTCT	660
	661	CAGCATGTACGAGGAAGGAAGTACTGCGCTAAAGGCCCTCCCACTCACCAAGGAGCTATT	720
	721	GGCTATTGATGATTGCTGAGGGAAGGGAGTAATTTTTTTTCTCTTTCTGAAGTGTGACTT	780
40	781	GAGTAAATTGCCCATAGTTCAGTATATAATCCCCAACCTGTGCTCAGGCAAGCAACCCTA	840
	841	ATTAAATGCAATAGCCACATACAAAAGAAGAGGATATGAATAGTTTGGTAGGAGGGGCTT	900
45	901	GTTAGGAAGAAGACATTAACAGGAGAGAGAGGAGCGAGAGGATAGTGAGTGTGTGAGAGT	960
	961	GCCTGCACGTGTGAAATGGTCAAAGAATTAAAAAATAAAAACTTAAAAAGCTATTAAAAA	1020
50	1021	GTAAAAAAATAAAA 1034	

Table 1 (continued):

human Teck cDNA (see SEQ ID NO: 3); signal sequence cleavage is probably between about Thr and Gln. Hu TECK protein sequence (see SEQ ID NO: 4).

5	TCGACCCACG CGTCCGCTTG GCCTACAGCC CGGCGGGCAT CAGCTCCCTT GACCCAGTGG	60
	ATATCGGTGG CCCCCTTATT CGTCCAGGTG CCCAGGGAGG AGGACCCGCC TGCAGC	116
10	ATG AAC CTG TGG CTC CTG GCC TGC CTG GTG GCC GGC TTC CTG GGA GCC Met Asn Leu Trp Leu Leu Ala Cys Leu Val Ala Gly Phe Leu Gly Ala -23 -20 -15 -10	164
15	TGG GCC CCC GCT GTC CAC ACC CAA GGT GTC TTT GAG GAC TGC TGC CTG Trp Ala Pro Ala Val His Thr Gln Gly Val Phe Glu Asp Cys Cys Leu -5 1 5	212
20	GCC TAC CAC TAC CCC ATT GGG TGG GCT GTG CTC CGG CGC GCC TGG ACT Ala Tyr His Tyr Pro Ile Gly Trp Ala Val Leu Arg Arg Ala Trp Thr 10 15 20 25	260
	TAC CGG ATC CAG GAG GTG AGC GGG AGC TGC AAT CTG CCT GCT GCG ATA Tyr Arg Ile Gln Glu Val Ser Gly Ser Cys Asn Leu Pro Ala Ala Ile 30 35 40	308
25	TTC TAC CTC CCC AAG AGA CAC AGG AAG GTG TGT GGG AAC CCC AAA AGC Phe Tyr Leu Pro Lys Arg His Arg Lys Val Cys Gly Asn Pro Lys Ser 45 50 55	356
30	AGG GAG GTG CAG AGA GCC ATG AAG CTC CTG GAT GCT CGA AAT AAG GTT Arg Glu Val Gln Arg Ala Met Lys Leu Leu Asp Ala Arg Asn Lys Val 60 65 70	404
35	TTT GCA AAG CTC CAC CAC AAC ATG CAG ACC TTC CAA GCA GGC CCT CAT Phe Ala Lys Leu His His Asn Met Gln Thr Phe Gln Ala Gly Pro His 75 80 85	452
40	GCT GTA AAG AAG TTG AGT TCT GGA AAC TCC AAG TTA TCA TCA TCC AAG Ala Val Lys Lys Leu Ser Ser Gly Asn Ser Lys Leu Ser Ser Ser Lys 90 95 100 105	500
	TTT AGC AAT CCC ATC AGC AGC AGC AAG AGG AAT GTC TCC CTC CTG ATA Phe Ser Asn Pro Ile Ser Ser Ser Lys Arg Asn Val Ser Leu Leu Ile 110 115 120	548
45	TCA GCT AAT TCA GGA CTG TGAGCCGGCT CATTTCTGGG CTCCATCGGC Ser Ala Asn Ser Gly Leu 125	596
50	ACAGGAGGGG CCGGATCTTT CTCCGATAAA ACCGTCGCCC TACAGACCCA GCTGTCCCCA	656
	CGCCTCTGTC TTTTGGGTCA AGTCTTAATC CCTGCACCTG AGTTGGTCCT CCCTCTGCAC	716
55	CCCCACCACC TCCTGCCCCGT CTGGCAACTG GAAAGAAGGA GTTGGCCTGA TTTTAACCTT	776
	TTGCCGCTCC GGGGAACAGC ACAATCCTGG GCAGCCAGTG GCTCTGTAG AGAAACTTA	836
	GGATACCTCT CTCACCTTCT GTTTCTTGCC GTCCACCCCG GGCCATGCCA GTGTGTCCCTC	896

Table 1 (continued):

TGGGTCCCCT CCAAAAATCT GGTCATTCAA GGATCCCCTC CCAAGGCTAT GCTTTTCTAT 956  
 5 AACTTTTAAA TAAACCTTGG GGGGTGAATG GAATAAAAAA AAAAAAAAAA AAAAAA 1012

10 Table 2: Nucleotide sequence (5' to 3') of MIP-3 $\alpha$  from human and the corresponding amino acid sequence (amino to carboxy), see SEQ ID NO: 5 and 6 and GenBank Accession U77035.

ATG TGC TGT ACC AAG AGT TTG CTC CTG GCT GCT TTG ATG TCA GTG CTG 48  
 Met Cys Cys Thr Lys Ser Leu Leu Leu Ala Ala Leu Met Ser Val Leu  
 15 -26 -25 -20 -15

CTA CTC CAC CTC TGC GGC GAA TCA GAA GCA GCA AGC AAC TTT GAC TGC 96  
 Leu Leu His Leu Cys Gly Glu Ser Glu Ala Ala Ser Asn Phe Asp Cys  
 -10 -5 1 5

20 TGT CTT GGA TAC ACA GAC CGT ATT CTT CAT CCT AAA TTT ATT GTG GGC 144  
 Cys Leu Gly Tyr Thr Asp Arg Ile Leu His Pro Lys Phe Ile Val Gly  
 10 15 20

25 TTC ACA CGG CAG CTG GCC AAT GAA GGC TGT GAC ATC AAT GCT ATC ATC 192  
 Phe Thr Arg Gln Leu Ala Asn Glu Gly Cys Asp Ile Asn Ala Ile Ile  
 25 30 35

30 TTT CAC ACA AAG AAA AAG TTG TCT GTG TGC GCA AAT CCA AAA CAG ACT 240  
 Phe His Thr Lys Lys Lys Leu Ser Val Cys Ala Asn Pro Lys Gln Thr  
 40 45 50

TGG GTG AAA TAT ATT GTG CGT CTC CTC AGT AAA AAA GTC AAG AAC ATG 288  
 Trp Val Lys Tyr Ile Val Arg Leu Leu Ser Lys Lys Val Lys Asn Met  
 35 55 60 65 70

TAAAACTGT GGCTTTTCTG GAATGGAATT GGACATAGCC CAAGAACAGA AAGAACCTTG 348

40 CTGGGGTTGG AGGTTTCACT TGCACATCAT GGAGGGTTTA GTGCTTATCT AATTTGTGCC 408

TCACTGGACT TGTCCAATTA ATGAAGTTGA TTCATATTGC ATCATAGTTT GCTTTGTTTA 468

AGCATCACAT TAAAGTTAAA CTGTATTTTA TGTTATTTAT AGCTGTAGGT TTTCTGTGTT 528

45 TAGCTATTTA ATACTAATTT TCCATAAGCT ATTTTGTTTT AGTGCAAAGT ATAAAATTAT 588

ATTTGGGGGG GAATAAGATT ATATGGACTT TTTTGCAAGC AACAAGCTAT TTTTAAAAA 648

50 AAACTATTTA ACATTCTTTT GTTTATATTG TTTTGTCTCC TAAATTGTTG TAATTGCATT 708

ATAAAATAAG AAAAATATTA ATAAGACAAA TATTGAAAAT AAAGAAACAA AAAGTTAAAA 768

AAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAA 801

Table 3: Nucleotide sequence (5' to 3') of MIP-3 $\beta$  from human and the corresponding amino acid sequence (amino to carboxy), see SEQ ID NO: 7 and 8, and GenBank Accession U77180. Signal sequence cleavage is about between Ser and Gly.

5	1	GGCACGAGCGGCACGAGCATCACTCACACCTTGCATTTACCCCTGCATCCCAGTCGCCC	60
	61	TGCAGCCTCACACAGATCCTGCACACACCCAGACAGCTGGCGCTCACACATTCACCGTTG	120
10	121	GCCTGCCTCTGTTTACCCTCCATGGCCCTGCTACTGGCCCTCAGCCTGCTGGTTCTCTGG	180
	1	MetAlaLeuLeuLeuAlaLeuSerLeuLeuValLeuTrp	13
	181	ACTTCCCCAGCCCCAACTCTGAGTGGCACCAATGATGCTGAAGACTGCTGCCTGTCTGTG	240
15	14	ThrSerProAlaProThrLeuSerGlyThrAsnAspAlaGluAspCysCysLeuSerVal	33
	241	ACCCAGAAACCCATCCCTGGGTACATCGTGAGGAACCTTCCACTACCTTCTCATCAAGGAT	300
	34	ThrGlnLysProIleProGlyTyrIleValArgAsnPheHisTyrLeuLeuIleLysAsp	53
	301	GGCTGCAGGGTGCCTGCTGTAGTGTTCACCACACTGAGGGGCCGCCAGCTCTGTGCACCC	360
20	54	GlyCysArgValProAlaValValPheThrThrLeuArgGlyArgGlnLeuCysAlaPro	73
	361	CCAGACCAGCCCTGGGTAGAACGCATCATCCAGAGACTGCAGAGGACCTCAGCCAAGATG	420
	74	ProAspGlnProTrpValGluArgIleIleGlnArgLeuGlnArgThrSerAlaLysMet	93
25	421	AAGCGCCGCAGCAGTTAACCTATGACCGTGCAGAGGGAGCCCGGAGTCCGAGTCAAGCAT	480
	94	LysArgArgSerSerEnd	98
	481	TGTGAATTATTACCTAACCTGGGGAACCGAGGACCAGAAGGAAGGACCAGGCTTCCAGCT	540
30	541	CCTCTGCACCAGACCTGACCAGCCAGGACAGGGCCTGGGGTGTGTGTGAGTGTGAGTGTG	600
	601	AGCGAGAGGGTGAGTGTGGTCTAGAGTAAAGCTGCTCCACCCCCAGATTGCAATGCTACC	660
35	661	AATAAAGCCGCCTGGTGTTTACAATAAAAAAAAAAAAAA	699

Table 4: Nucleotide sequence (5' to 3') of chemokine receptor, DC CR, from human and the corresponding amino acid sequence (amino to carboxy), see SEQ ID NO: 9 and 10. Nucleotide 579 may be A, C, G, or T, and the codon may code for His or Gln.

5	1	ATGTTTTCGACTCCAGTGAAGATTATTTTGTGTGTCAGTCAATACTTCATATTACTCAGTTG	60
	1	MetPheSerThrProValLysIleIleLeuCysGlnSerIleLeuHisIleThrGlnLeu	20
	61	ATTCTGAGATGTTACTGTGCTCCTTGCAGGAGGTCAGGCAGTTCTCCAGGCTATTTGTAC	120
10	21	IleLeuArgCysTyrCysAlaProCysArgArgSerGlySerSerProGlyTyrLeuTyr	40
	121	CGAATTGCCTACTCCTTGATCTGTGTTCTTGGCCTCCTGGGGAATATTCTGGTGGTGATC	180
	41	ArgIleAlaTyrSerLeuIleCysValLeuGlyLeuLeuGlyAsnIleLeuValValIle	60
15	181	ACCTTTGCTTTTTTATAAGAAGGCCAGGTCTATGACAGACGTCCTATCTCTGAACATGGCC	240
	61	ThrPheAlaPheTyrLysLysAlaArgSerMetThrAspValTyrLeuLeuAsnMetAla	80
	241	ATTGCAGACATCCTCTTTGTTCTTACTCTCCCATTTCTGGGCAGTGAGTCATGCCACTGGT	300
20	81	IleAlaAspIleLeuPheValLeuThrLeuProPheTrpAlaValSerHisAlaThrGly	100
	301	GCGTGGGTTTTTCAGCAATGCCACGTGCAAGTTGCTAAAAGGCATCTATGCCATCAACTTT	360
	101	AlaTrpValPheSerAsnAlaThrCysLysLeuLeuLysGlyIleTyrAlaIleAsnPhe	120
25	361	AACTGCGGGATGCTGCTCCTGACTTGCATTAGCATGGACCGGTACATCGCCATTGTACAG	420
	121	AsnCysGlyMetLeuLeuLeuThrCysIleSerMetAspArgTyrIleAlaIleValGln	140
	421	GCGACTAAGTCATTCCGGCTCCGATCCAGAACACTACCGCGCAGCAAATCATCTGCCTT	480
	141	AlaThrLysSerPheArgLeuArgSerArgThrLeuProArgSerLysIleIleCysLeu	160
30	481	GTTGTGTGGGGGCTGTGTCAGTCATCATCTCCAGCTCAACTTTTGTCTTCAACCAAAAATAC	540
	161	ValValTrpGlyLeuSerValIleIleSerSerSerThrPheValPheAsnGlnLysTyr	180
	541	AACACCCAAGGCAGCGATGTCTGTGAACCCAAGTACCAAACTGTCTCGGAGCCCATCAGG	600
35	181	AsnThrGlnGlySerAspValCysGluProLysTyrGlnThrValSerGluProIleArg	200
	601	TGGAAGCTGCTGATGTTGGGGCTTGAGCTACTCTTTGGTTTCTTTATCCCTTTGATGTTT	660
	201	TrpLysLeuLeuMetLeuGlyLeuGluLeuLeuPheGlyPhePheIleProLeuMetPhe	220
40	661	ATGATATTTTGTACACGTTTATTGTCAAACCTTGGTGCAAGCTCAGAATTCTAAAAGG	720
	221	MetIlePheCysTyrThrPheIleValLysThrLeuValGlnAlaGlnAsnSerLysArg	240
	721	CACAAAGCCATCCGTGTAATCATAGCTGTGGTGCTTGTGTTTCTGGCTTGTGAGATTCT	780
	241	HisLysAlaIleArgValIleIleAlaValValLeuValPheLeuAlaCysGlnIlePro	260
45	781	CATAACATGGTCCTGCTTGTGACGGCTGCTAATTTGGGTAAAATGAACCGATCCTGCCAG	840
	261	HisAsnMetValLeuLeuValThrAlaAlaAsnLeuGlyLysMetAsnArgSerCysGln	280
	841	AGCGAAAAGCTAATTGGCTATACGAAAACGTGTCACAGAAGTCCTGGCTTTCTCTGCACTGC	900
50	281	SerGluLysLeuIleGlyTyrThrLysThrValThrGluValLeuAlaPheLeuHisCys	300
	901	TGCCTGAACCTGTGCTCTACGCTTTTATTGGGCAGAAGTTCAGAACTACTTTCTGAAG	960
	301	CysLeuAsnProValLeuTyrAlaPheIleGlyGlnLysPheArgAsnTyrPheLeuLys	320
55	961	ATCTTGAAGGACCTGTGGTGTGTGAGAAGGAAGTACAAGTCCTCAGGCTTCTCTGTGCC	1020
	321	IleLeuLysAspLeuTrpCysValArgArgLysTyrLysSerSerGlyPheSerCysAla	340
	1021	GGGAGGTACTCAGAAAACATTTCTCGGCAGACCAGTGAGACCGCAGATAACGACAATGCG	1080
	341	GlyArgTyrSerGluAsnIleSerArgGlnThrSerGluThrAlaAspAsnAspAsnAla	360

Table 4 (continued)

1081 TCGTCCTTCACTATGTGATAGAAAGCTGAGTCTCCCTAA 1119  
 361 SerSerPheThrMetEnd 365

5

Table 5: Nucleotide sequence (5' to 3') of chemokine receptor, M/DC CR, from human and the corresponding amino acid sequence (amino to carboxy), see SEQ ID NO: 11 and 12.

10

1	GAGGAAGCTGCTTCGGGGGGTGAGCAAACCTTTTAAATGCAGAAATTATGATCTACACC	60
	MetIleTyrThr	4
61	CGTTTCTTAAAAGGCAGTCTGAAGATGGCCAATTACACGCTGGCACCAGAGGATGAATAT	120
15	5 ArgPheLeuLysGlySerLeuLysMetAlaAsnTyrThrLeuAlaProGluAspGluTyr	24
121	GATGTCCTCATAGAAGGTGAAGTGGAGAGCGATGAGGCAGAGCAATGTGACAAGTATGAC	180
25	AspValLeuIleGluGlyGluLeuGluSerAspGluAlaGluGlnCysAspLysTyrAsp	44
181	GCCCAGGCACTCTCAGCCCAGCTGGTGCCATCACTCTGCTCTGCTGTGTTTGTGATCGGT	240
20	45 AlaGlnAlaLeuSerAlaGlnLeuValProSerLeuCysSerAlaValPheValIleGly	64
241	GTCCTGGACAATCTCCTGGTTGTGCTTATCCTGGTAAATATAAAGGACTCAAACGCGTG	300
25	65 ValLeuAspAsnLeuLeuValValLeuIleLeuValLysTyrLysGlyLeuLysArgVal	84
301	GAAAATATCTATCTTCTAAACTTGGCAGTTTCTAACTTGTGTTTCTTGCTTACCCTGCCC	360
85	GluAsnIleTyrLeuLeuAsnLeuAlaValSerAsnLeuCysPheLeuLeuThrLeuPro	104
361	TTCTGGGCTCATGCTGGGGGCGATCCCATGTGTAAATTTCTCATTGGACTGTACTTCGTG	420
30	105 PheTrpAlaHisAlaGlyGlyAspProMetCysLysIleLeuIleGlyLeuTyrPheVal	124
421	GGCCTGTACAGTGAGACATTTTCAATTGCCTTCTGACTGTGCAAAGGTACCTAGTGTTT	480
125	GlyLeuTyrSerGluThrPhePheAsnCysLeuLeuThrValGlnArgTyrLeuValPhe	144
481	TTGCACAAGGGCAACTTTTCTCAGCCAGGAGGAGGGTGCCCTGTGGCATCATTACAAGT	540
35	145 LeuHisLysGlyAsnPhePheSerAlaArgArgArgValProCysGlyIleIleThrSer	164
541	GTCCTGGCATGGGTAACAGCCATTCTGGCCACTTTGCCTGAATTCGTGGTTTATAAACCT	600
40	165 ValLeuAlaTrpValThrAlaIleLeuAlaThrLeuProGluPheValValTyrLysPro	184
601	CAGATGGAAGACCAGAAATACAAGTGTGCATTTAGCAGAACTCCCTTCCTGCCAGCTGAT	660
185	GlnMetGluAspGlnLysTyrLysCysAlaPheSerArgThrProPheLeuProAlaAsp	204
661	GAGACATTCTGGAAGCATTTTCTGACTTTAAAAATGAACATTTCCGGTTCTTGTCCTCCCC	720
45	205 GluThrPheTrpLysHisPheLeuThrLeuLysMetAsnIleSerValLeuValLeuPro	224
721	CTATTTATTTTACATTTCTCTATGTGCAAATGAGAAAAACACTAAGGTTTCAGGGAGCAG	780
225	LeuPheIlePheThrPheLeuTyrValGlnMetArgLysThrLeuArgPheArgGluGln	244
781	AGGTATAGCCTTTTCAAGCTTGTTTTTGCCGTAATGGTAGTCTTCCTTCTGATGTGGGCG	840
50	245 ArgTyrSerLeuPheLysLeuValPheAlaValMetValValPheLeuLeuMetTrpAla	264
841	CCCTACAATATTGCATTTTTCCTGTCCACTTTCAAAGAACACTTCTCCCTGAGTGACTGC	900
55	265 ProTyrAsnIleAlaPhePheLeuSerThrPheLysGluHisPheSerLeuSerAspCys	284
901	AAGAGCAGCTACAATCTGGACAAAAGTGTTTCACATCACTAAACTCATCGCCACCACCCAC	960
285	LysSerSerTyrAsnLeuAspLysSerValHisIleThrLysLeuIleAlaThrThrHis	304
961	TGCTGCATCAACCCTCTCCTGTATGCGTTTCTTGATGGGACATTTAGCAAATACCTCTGC	1020
60	305 CysCysIleAsnProLeuLeuTyrAlaPheLeuAspGlyThrPheSerLysTyrLeuCys	324

1021	CGCTGTTTCCATCTGCGTAGTAACACCCCACTTCAACCCAGGGGGCAGTCTGCACAAGGC	1080
325	ArgCysPheHisLeuArgSerAsnThrProLeuGlnProArgGlyGlnSerAlaGlnGly	344
1081	ACATCGAGGGAAGAACCTGACCATTCCACCGAAGTGTAACCTAGCATCCACCAAAATGCAA	1140
345	ThrSerArgGluGluProAspHisSerThrGluValEnd	356
1141	GAAGAATAAACATGGATTTTTCATCTTTCTGCATTATTTTCATGTAAATTTTCTACACATTT	1200
1201	GTATACAAAATCGGATACAGGAAGAAAAGGGAGAGGTGAGCTAACATTTGCTAAGCACTG	1260
1261	AATTTGTCTCAGGCACCGTGCAAGGCTCTTTACAAACGTGAGCTCCTTCGCCTCCTACCA	1320
1321	CTTGTCCATAGTGTGGATAGGACTAGTCTCATTTCTCTGAGAAGAAAATAAGGCGCGGA	1380
1381	AATTTGTCTAAGATCACATAACTAGGAAGTGGCAGAACTGATTCTCCAGCCCTGGTAGCA	1440
1441	TTTGCTCAGAGCCTACGCTTGGTCCAGAACATCAAAC TCCAAACCCTGGGGACAAACGAC	1500
1501	ATGAAAATAAATGTATTTTAAACATATAAAAAAAAAAAAAAAAAAAAAA	1547

```

30 M/DC CR      MIYTRFLKGS LKMAN YTLAPEDEYD VLI EGEL ESDEAE QCDKYDAQALS
   C-C CKR-1      METPNTTEDY DTTTEFDYGDATPCQKVNERAFG
   C-C CKR-2      MLSTSRSRFIRNTNESGEEVTTFFDYDYGAPCHKFDVKQIG
   C-C CKR-3      MTTSLD TVETFGTTSY YDDVGLLC EKADTRALM
   C-C CKR-4      MNPTDIAD TTLDES IYSNYYLYESIPKPCTKEGIKAFG
35

```

40 M/DC CR AQLVPSLCSAVFVIGVLDNLLVVLILVKYKGLKRVENIYLLNLAVSNLCF  
C-C CKR-1 AQLLPPLYSLVVFVIGLVGNILVVLVLVQYKRLKNMTSIYLLNLAI SDLLF  
C-C CKR-2 AQLLPPLYSLVFIFGFVGNMLVVLILINCKKLKCLTDIYLLNLAI SDLLF  
C-C CKR-3 AQFVPPLYSLVFTVGLLGNVVVMILIKYRRLRIMTNIYLLNLAI SDLLF  
C-C CKR-4 ELFLPPLYSLVVFVGLLGNVSVVLVLFKYKRLRSMTDVYLLNLAI SDLLF  
\* \* \* \* \*

```

M/DC CR      LLTLPFWAHAG-----GDPMKILIGLYFVGLYSETFFNCLLTVQRYL
45 C-C CKR-1   LFTLPFWIDYKLKDDWVF GDAMCKILSGFYTGlySEIFFIILLTIDRYL
C-C CKR-2     LITLPLWAH-SANEWVF GNAMCKLFTGLYHIGYFGGIFFIILLTIDRYL
C-C CKR-3     LVTLFPWIHYVRGHNVWF GHGMCKLLSGFYHTGLYSEIFFIILLTIDRYL
C-C CKR-4     VFSLPFWGYGA-ADQWVFLGLCKMISWMYLVGfySGIFFVMlmsIDRYL
          .  *  *  *                *    *  *  *            *    *  *  *  *  *  *

```

```

55 M/DC CR      VFLHKGNNFFSAR-RRVPCGIITSVLAWVTAILATLPEFVVYKPMEDQKY
   C-C CKR-1   AIVH--AVFALRARTVTFGVITSIIIWALAILASMPGLYFSKTQWEFTHH
   C-C CKR-2   AIVH--AVFALKARTVTFGVVTSVITWLVAVFASVPGIIFTKCQKEDSVY
   C-C CKR-3   AIVH--AVFALRARTVTFGVITSIVTWGLAVLAALPEFIFYETEELFEET
   C-C CKR-4   AIVH--AVFSLRARTLTGYVITSLATWSVAVFASLPGFLFSTCYTERNHT
              . * . * . * . * . * . * . * . * .

```



Table 6 (continued)

5	M/DC CR	KCAFSRTPFLPADETF-WKHFLTLKMNISVLVLPFLFIFTFLYVQMRKTL-
	C-C CKR-1	TCS----LHFPHESLREWKLQALKLNLFGLVLPPLVMIICYTGIIKILL
	C-C CKR-2	VCG----PYFPR----GWNNFHTIMRNILGLVLPPLLMVICYSGILKTLL
	C-C CKR-3	LCS----ALYPEDTVYSWRHFHTLRMTIFCLVLPPLLVMAICYTGIIKTLL
	C-C CKR-4	YCK----TKYSLNST-TWKVLSSLEINILGLVIPLGIMLFCYSMIIRTLO
		* . . . . . ** . . . *
10	M/DC CR	--RFREQRYSLFKLVFAVMVVFLLMWAPYNIAFFLSTFKEHFSLSACKSS
	C-C CKR-1	RRPNEKK-SKAVRLIFVIMIIFFLFWTPYNLTILISVFQDFLFTHECEQS
	C-C CKR-2	RCRNEKKRHRAVRVIFTIMIVYFLFWTPYNIVILLNTFQEFFGLSNCEST
	C-C CKR-3	RCPSKKK-YKAIRLIFVIMAVFFIFWTPYNVAILLSSYQSILFGNDCERS
	C-C CKR-4	HCKNEKK-NKAVKMIFAVVFLGLFWTPYNIVLFLETFLVELEVLDCTFE
15		. . . * . . . . . * . . . . . *
20	M/DC CR	YNLDKSVHITKLIATTHCCINPLLYAFLDGTFSKYLRCRCFH-----
	C-C CKR-1	RHLDLAVQVTEVIAIYTHCCVNPVIYAFVGERFRKYLRQLFH-RRVA----
	C-C CKR-2	SQLDQATQVTETLGMTHCCINPIIYAFVGEKFRSLFHIALG-CRIAPLQK
	C-C CKR-3	KHLDLVMLVTEVIAIYSHCCMNPIYAFVGERFRKYLRHFFH-RHLL----
	C-C CKR-4	RYLDYAIQATETLAFVHCCLNPIIYFFLGEKFRKYILQLFKTCRGLFVLC
		** . . . . . * . . . . . * . . . . . *
25	M/DC CR	-----LRSNTPLQPRGQSAQGTSREEP--DHSTEV*
	C-C CKR-1	-----VHLVKWLPFLSVDRLERVSSTSPSTGEHELSA----GF*
	C-C CKR-2	PVCGGPGVRPGKNVKVTTQGLLDGRGKGKSIGRAPEASLQDKEGA*
	C-C CKR-3	-----MHLGRYIPFLPSEKLERTSSVSPSTAEPESL----VF*
	C-C CKR-4	QYCG-----LLQIYSAD-----TPSSSYTQSTMDHDLHDAL*

30

As used herein, the term "TECK" shall encompass, when used in a protein context, a protein having mature mouse or human amino acid sequences, as shown in Table 1. The invention also embraces a polypeptide comprising a significant fragment of such protein. It also refers to a polypeptide which is a species counterpart, e.g., which exhibits similar biological function, and is more homologous in natural encoding sequence than other genes from that species. Typically, such chemokine will also interact with its specific binding components, e.g., receptor. These binding components, e.g., antibodies, typically bind to the chemokine with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM. Homologous proteins would be found in mammalian species other than mouse, e.g., rats, dogs, cats, and primates. Non-mammalian species should also

45

possess structurally or functionally related genes and proteins.

The term "polypeptide" as used herein includes a significant fragment or segment, and encompasses a stretch  
5 of amino acid residues of at least about 8 amino acids, generally at least 10 amino acids, more generally at least 12 amino acids, often at least 14 amino acids, more often at least 16 amino acids, typically at least 18 amino acids, more typically at least 20 amino acids, usually at least 22  
10 amino acids, more usually at least 24 amino acids, preferably at least 26 amino acids, more preferably at least 28 amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids, e.g., about 35, 40, 45, 50, 60, 75, 80, 100, 120, etc. Similar  
15 proteins will likely comprise a plurality of such segments. Such fragments may have ends which begin and/or end at virtually all positions, e.g., beginning at residues 1, 2, 3, etc., and ending at, e.g., 69, 68, 67, 66, etc., in all combinations. Particularly interesting peptides have ends  
20 corresponding to structural domain boundaries. See, e.g., PHD and DSC programs, Rost and Sander (1994) Proteins 19:55-72; and King and Sternberg (1996) Protein Science 5:2298-2310.

The term "binding composition" refers to molecules  
25 that bind with specificity to the respective chemokine or receptor, e.g., in a ligand-receptor type fashion or an antibody-antigen interaction. These compositions may be compounds, e.g., proteins, which specifically associate with the chemokine or receptor, including natural  
30 physiologically relevant protein-protein interactions, either covalent or non-covalent. The binding composition may be a polymer, or another chemical reagent. No implication as to whether the chemokine presents a concave or convex shape in its ligand-receptor interaction is  
35 represented, other than the interaction exhibit similar specificity, e.g., specific affinity. A functional analog may be a ligand with structural modifications, or may be a

wholly unrelated molecule, e.g., which has a molecular shape which interacts with the appropriate ligand binding determinants. The ligands may serve as agonists or antagonists of the receptor, see, e.g., Goodman, et al.

- 5 (eds.) (1990) Goodman & Gilman's: The Pharmacological Bases of Therapeutics (8th ed.), Pergamon Press.

Substantially pure means that the protein is free from other contaminating proteins, nucleic acids, and/or other biologicals typically derived from the original source  
10 organism. Purity may be assayed by standard methods, and will ordinarily be at least about 40% pure, more ordinarily at least about 50% pure, generally at least about 60% pure, more generally at least about 70% pure, often at least about 75% pure, more often at least about 80% pure,  
15 typically at least about 85% pure, more typically at least about 90% pure, preferably at least about 95% pure, more preferably at least about 98% pure, and in most preferred embodiments, at least 99% pure. Analyses will typically be by weight, but may be by molar amounts.

20 Solubility of a polypeptide or fragment depends upon the environment and the polypeptide. Many parameters affect polypeptide solubility, including temperature, electrolyte environment, size and molecular characteristics of the polypeptide, and nature of the solvent. Typically,  
25 the temperature at which the polypeptide is used ranges from about 4° C to about 65° C. Usually the temperature at use is greater than about 18° C and more usually greater than about 22° C. For diagnostic purposes, the temperature will usually be about room temperature or warmer, but less  
30 than the denaturation temperature of components in the assay. For therapeutic purposes, the temperature will usually be body temperature, typically about 37° C for humans, though under certain situations the temperature may be raised or lowered in situ or in vitro.

35 The electrolytes will usually approximate in situ physiological conditions, but may be modified to higher or lower ionic strength where advantageous. The actual ions

may be modified, e.g., to conform to standard buffers used in physiological or analytical contexts.

The size and structure of the polypeptide should generally be in a substantially stable state, and usually not in a denatured state. The polypeptide may be associated with other polypeptides in a quaternary structure, e.g., to confer solubility, or associated with lipids or detergents in a manner which approximates natural lipid bilayer interactions.

The solvent will usually be a biologically compatible buffer, of a type used for preservation of biological activities, and will usually approximate a physiological solvent. Usually the solvent will have a neutral pH, typically between about 5 and 10, and preferably about 7.5. On some occasions, a detergent will be added, typically a mild non-denaturing one, e.g., CHS or CHAPS, or a low enough concentration as to avoid significant disruption of structural or physiological properties of the protein.

Solubility is reflected by sedimentation measured in Svedberg units, which are a measure of the sedimentation velocity of a molecule under particular conditions. The determination of the sedimentation velocity was classically performed in an analytical ultracentrifuge, but is typically now performed in a standard ultracentrifuge. See, Freifelder (1982) Physical Biochemistry (2d ed.), W.H. Freeman; and Cantor and Schimmel (1980) Biophysical Chemistry, parts 1-3, W.H. Freeman & Co., San Francisco. As a crude determination, a sample containing a putatively soluble polypeptide is spun in a standard full sized ultracentrifuge at about 50K rpm for about 10 minutes, and soluble molecules will remain in the supernatant. A soluble particle or polypeptide will typically be less than about 30S, more typically less than about 15S, usually less than about 10S, more usually less than about 6S, and, in particular embodiments, preferably less than about 4S, and more preferably less than about 3S.

### III. Physical Variants

This invention also encompasses proteins or peptides having substantial amino acid sequence homology with the amino acid sequence of each respective chemokine or receptor. The variants include species or polymorphic variants.

Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches, if necessary, by introducing gaps as required. This changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences are typically intended to include natural allelic and interspecies variations in each respective protein sequence. Typical homologous proteins or peptides will have from 25-100% homology (if gaps can be introduced), to 50-100% homology (if conservative substitutions are included) with the amino acid sequence of the appropriate chemokine or receptor. Homology measures will be at least about 35%, generally at least 40%, more generally at least 45%, often at least 50%, more often at least 55%, typically at least 60%, more typically at least 65%, usually at least 70%, more usually at least 75%, preferably at least 80%, and more preferably at least 80%, and in particularly preferred embodiments, at least 85% or more. See also Needleham, et al. (1970) J. Mol. Biol. 48:443-453; Sankoff, et al. (1983) Chapter One in Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics Computer Group, Madison, WI.

Each of the isolated chemokine or receptor DNAs can be readily modified by nucleotide substitutions, nucleotide deletions, nucleotide insertions, and inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode these antigens, their derivatives, or proteins having similar physiological, immunogenic, or antigenic activity. These modified sequences can be used to produce mutant antigens or to enhance expression. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant chemokine or receptor derivatives include predetermined or site-specific mutations of the respective protein or its fragments. "Mutant chemokine" encompasses a polypeptide otherwise falling within the homology definition of the chemokine as set forth above, but having an amino acid sequence which differs from that of the chemokine as found in nature, whether by way of deletion, substitution, or insertion. These include substitution levels from none, one, two, three, etc. In particular, "site specific mutant chemokine" generally includes proteins having significant homology with a ligand having sequences of Table 1 through 3, and as sharing various biological activities, e.g., antigenic or immunogenic, with those sequences, and in preferred embodiments contain most of the disclosed sequences. Similar concepts apply to the different chemokine protein embodiments, particularly those found in various warm blooded animals, e.g., mammals and birds. As stated before, it is emphasized that descriptions are generally meant to encompass the various chemokine proteins, not limited to the mouse or human embodiments specifically discussed. Similar concepts apply to the receptor embodiments.

Although site specific mutation sites are often predetermined, mutants need not be site specific. Chemokine mutagenesis can be conducted by making amino acid insertions or deletions. Substitutions, deletions,

insertions, or combinations may be generated to arrive at a final construct. Insertions include amino- or carboxy-terminal fusions. Random mutagenesis can be conducted at a target codon and the expressed mutants can then be screened  
5 for the desired activity. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis or polymerase chain reaction (PCR) techniques. See also Sambrook, et al. (1989) and Ausubel, et al. (1987  
10 and Supplements).

The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that could hybridize to produce secondary mRNA structure such as loops or hairpins.

15 The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using segments from these proteins. A heterologous fusion protein is a fusion of proteins or segments which are naturally not normally fused in the same manner. Thus, the fusion  
20 product of an immunoglobulin with a chemokine or receptor polypeptide is a continuous protein molecule having sequences fused in a typical peptide linkage, typically made as a single translation product and exhibiting properties derived from each source peptide. A similar  
25 chimeric concept applies to heterologous nucleic acid sequences.

In addition, new constructs may be made from combining similar functional domains from other proteins. For example, ligand-binding or other segments may be "swapped"  
30 between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of  
35 ligand-binding specificities and other functional domains.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce

suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence, e.g., PCR techniques.

#### IV. Functional Variants

The blocking of physiological response to various embodiments of these chemokines may result from the inhibition of binding of the ligand to its receptor, likely through competitive inhibition. Thus, in vitro assays of the present invention will often use isolated protein, membranes from cells expressing a recombinant membrane associated chemokine, soluble fragments comprising receptor binding segments of these ligands, or fragments attached to solid phase substrates. These assays will also allow for the diagnostic determination of the effects of either binding segment mutations and modifications, or ligand mutations and modifications, e.g., ligand analogs.

This invention also contemplates the use of competitive drug screening assays, e.g., where neutralizing antibodies to antigen or receptor fragments compete with a test compound for binding to the protein. In this manner, the antibodies can be used to detect the presence of polypeptides which share one or more antigenic binding sites of the ligand and can also be used to occupy binding sites on the protein that might otherwise interact with a receptor.

Additionally, neutralizing antibodies against a specific chemokine embodiment and soluble fragments of the chemokine which contain a high affinity receptor binding site, can be used to inhibit chemokine activity in tissues, e.g., tissues experiencing abnormal physiology.

"Derivatives" of chemokine antigens include amino acid sequence mutants, glycosylation variants, and covalent or aggregate conjugates with other chemical moieties.



Covalent derivatives can be prepared by linkage of functionalities to groups which are found in chemokine amino acid side chains or at the N- or C- termini, by means which are well known in the art. These derivatives can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl side chains, O-acyl derivatives of hydroxyl group-containing residues, and N-acyl derivatives of the amino terminal amino acid or amino-group containing residues, e.g., lysine or arginine. Acyl groups are selected from the group of alkyl-moieties including C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species. Covalent attachment to carrier proteins may be important when immunogenic moieties are haptens.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. Particularly preferred means for accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine.

A major group of derivatives are covalent conjugates of the respective chemokine or receptor or fragments thereof with other proteins or polypeptides. These derivatives can be synthesized in recombinant culture such as N- or C-terminal fusions or by the use of agents known in the art for their usefulness in cross-linking proteins through reactive side groups. Preferred chemokine derivatization sites with cross-linking agents are at free amino groups, carbohydrate moieties, and cysteine residues.

Fusion polypeptides between these chemokines and other homologous or heterologous proteins, e.g., other

chemokines, are also provided. Many growth factors and cytokines are homodimeric entities, and a repeat construct may have various advantages, including lessened susceptibility to proteolytic cleavage. Moreover, many  
5 receptors require dimerization to transduce a signal, and various dimeric ligands or domain repeats can be desirable. Homologous polypeptides may be fusions between different surface markers, resulting in, e.g., a hybrid protein exhibiting receptor binding specificity. Likewise,  
10 heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a ligand, e.g., a receptor-binding segment, so that the  
15 presence or location of the fused ligand may be easily determined. See, e.g., Dull, et al., U.S. Patent No. 4,859,609. Other gene fusion partners include bacterial  $\beta$ -galactosidase, trpE, Protein A,  $\beta$ -lactamase, alpha amylase, alcohol dehydrogenase, a FLAG fusion, and yeast alpha  
20 mating factor. See, e.g., Godowski, et al. (1988) Science 241:812-816.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded  
25 fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

30 Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation, sulfonation, biotinylation, or the addition or removal of other moieties, particularly those which have molecular shapes similar to phosphate groups. In some embodiments,  
35 the modifications will be useful labeling reagents, or serve as purification targets, e.g., affinity tags as FLAG.

Fusion proteins will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, for example, in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.), Vols. 1-3, Cold Spring Harbor Laboratory. Techniques for synthesis of polypeptides are described, for example, in Merrifield (1963) J. Amer. Chem. Soc. 85:2149-2156; Merrifield (1986) Science 232: 341-347; and Atherton, et al. (1989) Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford; and chemical ligation, e.g., Dawson, et al. (1994) Science 266:776-779, a method of linking long synthetic peptides by a peptide bond.

This invention also contemplates the use of derivatives of these chemokines or receptors other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives generally fall into the three classes: (1) salts, (2) side chain and terminal residue covalent modifications, and (3) adsorption complexes, for example with cell membranes. Such covalent or aggregative derivatives are useful as immunogens, as reagents in immunoassays, or in purification methods such as for affinity purification of ligands or other binding ligands. For example, a chemokine antigen can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated Sepharose, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of anti-chemokine antibodies or its receptor. These chemokines can also be labeled with a detectable group, for example radioiodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to a fluorescent moiety for use in diagnostic assays. Purification of chemokine may be effected by immobilized antibodies or receptor.

Other modifications may be introduced with the goal of modifying the therapeutic pharmacokinetics or pharmacodynamics of a target chemokine. For example, certain means to minimize the size of the entity may  
5 improve its pharmacoaccessibility; other means to maximize size may affect pharmacodynamics.

A solubilized chemokine or appropriate fragment of this invention can be used as an immunogen for the production of antisera or antibodies specific for the  
10 ligand or fragments thereof. The purified chemokines can be used to screen monoclonal antibodies or chemokine-binding fragments prepared by immunization with various forms of impure preparations containing the protein. In particular, antibody equivalents include antigen binding  
15 fragments of natural antibodies, e.g., Fv, Fab, or F(ab)<sub>2</sub>. Purified chemokines can also be used as a reagent to detect antibodies generated in response to the presence of elevated levels of the protein or cell fragments containing the protein, both of which may be diagnostic of an abnormal  
20 or specific physiological or disease condition. Additionally, chemokine protein fragments, or their concatenates, may also serve as immunogens to produce antibodies of the present invention, as described immediately below. For example, this invention  
25 contemplates antibodies raised against amino acid sequences shown in Tables 1 through 3, or proteins containing them. In particular, this invention contemplates antibodies having binding affinity to or being raised against specific fragments, e.g., those which are predicted to lie on the  
30 outside surfaces of protein tertiary structure. Similar concepts apply to antibodies specific for receptors of the invention.

The present invention contemplates the isolation of additional closely related species variants. Southern and  
35 Northern blot analysis should establish that similar genetic entities exist in other mammals, and establish the stringency of hybridization conditions to isolate such. It

is likely that these chemokines and receptors are widespread in species variants, e.g., rodents, lagomorphs, carnivores, artiodactyla, perissodactyla, and primates.

The invention also provides means to isolate a group  
5 of related chemokines displaying both distinctness and similarities in structure, expression, and function. Elucidation of many of the physiological effects of the proteins will be greatly accelerated by the isolation and characterization of distinct species variants of the  
10 ligands. In particular, the present invention provides useful probes for identifying additional homologous genetic entities in different species.

The isolated genes will allow transformation of cells lacking expression of a corresponding chemokine, e.g.,  
15 either species types or cells which lack corresponding ligands and exhibit negative background activity. Expression of transformed genes will allow isolation of antigenically pure cell lines, with defined or single specie variants. This approach will allow for more  
20 sensitive detection and discrimination of the physiological effects of chemokine receptor proteins. Subcellular fragments, e.g., cytoplasts or membrane fragments, can be isolated and used.

Dissection of critical structural elements which  
25 effect the various differentiation functions provided by ligands is possible using standard techniques of modern molecular biology, particularly in comparing members of the related class. See, e.g., the homolog-scanning mutagenesis technique described in Cunningham, et al. (1989) Science  
30 243:1339-1336; and approaches used in O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992; and Lechleiter, et al. (1990) EMBO J. 9:4381-4390.

In addition, receptor binding segments can be substituted between species variants to determine what  
35 structural features are important in both receptor binding affinity and specificity, as well as signal transduction. An array of different chemokine variants will be used to

screen for ligands exhibiting combined properties of interaction with different receptor species variants.

Intracellular functions would probably involve segments of the receptor which are normally accessible to the cytosol. However, ligand internalization may occur under certain circumstances, and interaction between intracellular components and "extracellular" segments may occur. The specific segments of interaction of a particular chemokine with other intracellular components may be identified by mutagenesis or direct biochemical means, e.g., cross-linking or affinity methods. Structural analysis by crystallographic or other physical methods will also be applicable. Further investigation of the mechanism of signal transduction will include study of associated components which may be isolatable by affinity methods or by genetic means, e.g., complementation analysis of mutants.

Further study of the expression and control of the various chemokines will be pursued. The controlling elements associated with the proteins may exhibit differential developmental, tissue specific, or other expression patterns. Upstream or downstream genetic regions, e.g., control elements, are of interest. Differential splicing of message may lead to membrane bound forms, soluble forms, and modified versions of ligand.

Structural studies of the proteins will lead to design of new ligands, particularly analogs exhibiting agonist or antagonist properties on the receptor. This can be combined with previously described screening methods to isolate ligands exhibiting desired spectra of activities.

Expression in other cell types will often result in glycosylation differences in a particular chemokine. Various species variants may exhibit distinct functions based upon structural differences other than amino acid sequence. Differential modifications may be responsible for differential function, and elucidation of the effects are now made possible.

Thus, the present invention provides important reagents related to a physiological chemokine-binding protein interaction. Although the foregoing description has focused primarily upon the mouse and human embodiments of the chemokines specifically described, those of skill in the art will immediately recognize that the invention provides other species counterparts, e.g., rat and other mammalian species or allelic or polymorphic variants.

## 10 V. Antibodies

Antibodies can be raised to these chemokines, including species or polymorphic variants, and fragments thereof, both in their naturally occurring forms and in their recombinant forms. Additionally, antibodies can be raised to chemokines in either their active forms or in their inactive forms. Anti-idiotypic antibodies are also contemplated.

Antibodies, including binding fragments and single chain versions, against predetermined fragments of the ligands can be raised by immunization of animals with concatemers or conjugates of the fragments with immunogenic proteins. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective chemokines, or screened for agonistic or antagonistic activity, e.g., mediated through a receptor for a chemokine. These monoclonal antibodies will usually bind with at least a  $K_D$  of about 1 mM, more usually at least about 300  $\mu$ M, typically at least about 10  $\mu$ M, more typically at least about 30  $\mu$ M, preferably at least about 10  $\mu$ M, and more preferably at least about 3  $\mu$ M or better.

The antibodies, including antigen binding fragments, of this invention can have significant diagnostic or therapeutic value. They can be potent antagonists that bind to ligand and inhibit binding to receptor or inhibit the ability of a ligand to elicit a biological response. They also can be useful as non-neutralizing antibodies and

can be coupled to toxins or radionuclides so that when the antibody binds to ligand, a cell expressing it, e.g., on its surface via receptor, is killed. Further, these antibodies can be conjugated to drugs or other therapeutic agents, either directly or indirectly by means of a linker, and may effect drug targeting. Antibodies to receptors may be more easily used to block ligand binding and signal transduction.

The antibodies of this invention can also be useful in diagnostic or reagent purification applications. As capture or non-neutralizing antibodies, they can be screened for ability to bind to the chemokines without inhibiting receptor binding. As neutralizing antibodies, they can be useful in competitive binding assays. They will also be useful in detecting or quantifying chemokine or, indirectly, receptors, e.g., in immunoassays. They may be used as purification reagents in immunoaffinity columns or as immunohistochemistry reagents.

Ligand fragments may be concatenated or joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. Short peptides will preferably be made as repeat structures to increase size. A ligand and its fragments may be fused or covalently linked to a variety of immunogens, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See Microbiology, Hoeber Medical Division, Harper and Row, 1969; Landsteiner (1962) Specificity of Serological Reactions, Dover Publications, New York, and Williams, et al. (1967) Methods in Immunology and Immunochemistry, Vol. 1, Academic Press, New York, for descriptions of methods of preparing polyclonal antisera. A typical method involves hyperimmunization of an animal with an antigen. The blood of the animal is then collected shortly after the repeated immunizations and the gamma globulin fraction is isolated.

In some instances, it is desirable to prepare monoclonal antibodies from various mammalian hosts, such as



mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) Basic and Clinical Immunology (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed.) Academic Press, New York; and particularly in Kohler and Milstein (1975) in Nature 256:495-497, which discusses one method of generating monoclonal antibodies. Summarized briefly, this method involves injecting an animal with an immunogen. The animal is then sacrificed and cells taken, e.g., from its spleen, which are then fused with myeloma cells. The result is a hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual clones, each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of immortalized and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic substance. Large amounts of antibody may be derived from ascites fluid from an animal.

Other suitable techniques involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," Science 246:1275-1281; and Ward, et al. (1989) Nature 341:544-546. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent

literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents, teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; and Queen et al. (1989) Proc. Nat'l. Acad. Sci. 86:10029-10033.

10       The antibodies of this invention can also be used for affinity chromatography in isolating the protein. Columns can be prepared where the antibodies are linked to a solid support, e.g., particles, such as agarose, Sephadex, or the like, where a cell lysate may be passed through the column, 15 the column washed, followed by increasing concentrations of a mild denaturant, whereby the purified chemokine protein will be released.

      The antibodies may also be used to screen expression libraries for particular expression products. Usually the 20 antibodies used in such a procedure will be labeled with a moiety allowing easy detection of presence of antigen by antibody binding.

      Antibodies raised against these chemokine will also be useful to raise anti-idiotypic antibodies. These will be 25 useful in detecting or diagnosing various immunological conditions related to expression of the respective antigens.

## VI. Nucleic Acids

30       The described peptide sequences and the related reagents are useful in isolating a DNA clone encoding these chemokines, e.g., from a natural source. Typically, it will be useful in isolating a gene from another individual, and similar procedures will be applied to isolate genes 35 from other species, e.g., warm blooded animals, such as birds and mammals. Cross hybridization will allow isolation of ligand from other species. A number of

different approaches should be available to successfully isolate a suitable nucleic acid clone. Similar concepts apply to the receptor embodiments.

The purified protein or defined peptides are useful  
5 for generating antibodies by standard methods, as described above. Synthetic peptides or purified protein can be presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) Current Protocols in Immunology Wiley/Greene; and Harlow and Lane  
10 (1989) Antibodies: A Laboratory Manual Cold Spring Harbor Press. Alternatively, a chemokine receptor can be used as a specific binding reagent, and advantage can be taken of its specificity of binding, much like an antibody would be used. However, chemokine receptors are typically 7  
15 transmembrane proteins, which could be sensitive to appropriate interaction with lipid or membrane. The signal transduction typically is mediated through a G-protein.

For example, the specific binding composition could be used for screening of an expression library made from a  
20 cell line which expresses a particular chemokine. The screening can be standard staining of surface expressed ligand, or by panning. Screening of intracellular expression can also be performed by various staining or immunofluorescence procedures. The binding compositions  
25 could be used to affinity purify or sort out cells expressing the ligand.

The peptide segments can also be used to predict appropriate oligonucleotides to screen a library, e.g., to isolate species variants. The genetic code can be used to  
30 select appropriate oligonucleotides useful as probes for screening. See, e.g., Tables 1 through 5. In combination with polymerase chain reaction (PCR) techniques, synthetic oligonucleotides will be useful in selecting correct clones from a library. Complementary sequences will also be used  
35 as probes or primers. Based upon identification of the likely amino terminus, the third peptide should be particularly useful, e.g., coupled with anchored vector or

poly-A complementary PCR techniques or with complementary DNA of other peptides.

This invention contemplates use of isolated DNA or fragments to encode a biologically active corresponding chemokine polypeptide. In addition, this invention covers isolated or recombinant DNA which encodes a biologically active protein or polypeptide which is capable of hybridizing under appropriate conditions with the DNA sequences described herein. Said biologically active protein or polypeptide can be an intact ligand, or fragment, and have an amino acid sequence as disclosed in Tables 1 through 3. Further, this invention covers the use of isolated or recombinant DNA, or fragments thereof, which encode proteins which are homologous to a chemokine or which was isolated using cDNA encoding a chemokine as a probe. The isolated DNA can have the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others. Alternatively, promoters or other regulatory signals may be incorporated to be operably linked to natural genes in a cell.

An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially separated from other components which naturally accompany a native sequence, e.g., ribosomes, polymerases, and flanking genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates and chemically synthesized analogs or analogs biologically synthesized by heterologous systems. A substantially pure molecule includes isolated forms of the molecule.

An isolated nucleic acid will generally be a homogeneous composition of molecules, but will, in some embodiments, contain minor heterogeneity. This heterogeneity is typically found at the polymer ends or

portions not critical to a desired biological function or activity.

A "recombinant" nucleic acid is defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the process is use of recombinant nucleic acid techniques, e.g., involving human intervention in the nucleotide sequence, typically selection or production. Alternatively, it can be a nucleic acid made by generating a sequence comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants. Thus, for example, products made by transforming cells with any unnaturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such is often done to replace a codon with a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a sequence recognition site. Alternatively, it is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms. Restriction enzyme recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode polypeptides similar to fragments of these antigens, and fusions of sequences from various different species variants.

A significant "fragment" in a nucleic acid context is a contiguous segment of at least about 17 nucleotides, generally at least about 20 nucleotides, more generally at least about 23 nucleotides, ordinarily at least about 26

nucleotides, more ordinarily at least about 29 nucleotides, often at least about 32 nucleotides, more often at least about 35 nucleotides, typically at least about 38 nucleotides, more typically at least about 41 nucleotides, usually at least about 44 nucleotides, more usually at least about 47 nucleotides, preferably at least about 50 nucleotides, more preferably at least about 53 nucleotides, and in particularly preferred embodiments will be at least about 56 or more nucleotides, e.g., 60, 65, 75, 85, 100, 120, 150, 200, 250, 300, 400, etc. Such fragments may have ends which begin and/or end at virtually all positions, e.g., beginning at nucleotides 1, 2, 3, etc., and ending at, e.g., 300, 299, 298, 287, etc., in all combinations. Particularly interesting polynucleotides have ends corresponding to structural domain boundaries.

A DNA which codes for a particular chemokine protein or peptide will be very useful to identify genes, mRNA, and cDNA species which code for related or homologous ligands, as well as DNAs which code for homologous proteins from different species. There are likely homologs in other species, including primates. Various chemokine proteins should be homologous and are encompassed herein. However, even proteins that have a more distant evolutionary relationship to the ligand can readily be isolated under appropriate conditions using these sequences if they are sufficiently homologous. Primate chemokines are of particular interest.

This invention further covers recombinant DNA molecules and fragments having a DNA sequence identical to or highly homologous to the isolated DNAs set forth herein. In particular, the sequences will often be operably linked to DNA segments which control transcription, translation, and DNA replication. Alternatively, recombinant clones derived from the genomic sequences, e.g., containing introns, will be useful for transgenic studies, including, e.g., transgenic cells and organisms, and for gene therapy. See, e.g., Goodnow (1992) "Transgenic Animals" in Roitt

- (ed.) Encyclopedia of Immunology Academic Press, San Diego, pp. 1502-1504; Travis (1992) Science 256:1392-1394; Kuhn, et al. (1991) Science 254:707-710; Capecchi (1989) Science 244:1288; Robertson (1987) (ed.) Teratocarcinomas and Embryonic Stem Cells: A Practical Approach IRL Press, Oxford; and Rosenberg (1992) J. Clinical Oncology 10:180-199.

Homologous nucleic acid sequences, when compared, exhibit significant similarity, or identity. The standards for homology in nucleic acids are either measures for homology generally used in the art by sequence comparison or based upon hybridization conditions. The hybridization conditions are described in greater detail below.

Substantial homology in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 50% of the nucleotides, generally at least about 56%, more generally at least about 59%, ordinarily at least about 62%, more ordinarily at least about 65%, often at least about 68%, more often at least about 71%, typically at least about 74%, more typically at least about 77%, usually at least about 80%, more usually at least about 85%, preferably at least about 90%, more preferably at least about 95 to 98% or more, and in particular embodiments, as high at about 99% or more of the nucleotides. Alternatively, substantial homology exists when the segments will hybridize under selective hybridization conditions, to a strand, or its complement, typically using a sequence derived from Tables 1 through 5. Typically, selective hybridization will occur when there is at least about 55% homology over a stretch of at least about 30 nucleotides, preferably at least about 65% over a stretch of at least about 25 nucleotides, more preferably at least about 75%, and most preferably at least about 90% over about 20 nucleotides. See, Kanehisa (1984) Nuc. Acids Res. 12:203-213. The length of homology comparison, as

described, may be over longer stretches, and in certain  
embodiments will be over a stretch of at least about 17  
nucleotides, usually at least about 20 nucleotides, more  
usually at least about 24 nucleotides, typically at least  
5 about 28 nucleotides, more typically at least about 40  
nucleotides, preferably at least about 50 nucleotides, and  
more preferably at least about 75 to 100 or more  
nucleotides.

Stringent conditions, in referring to homology in the  
10 hybridization context, will be stringent combined  
conditions of salt, temperature, organic solvents, and  
other parameters, typically those controlled in  
hybridization reactions. Stringent temperature conditions  
will usually include temperatures in excess of about 30° C,  
15 more usually in excess of about 37° C, typically in excess  
of about 45° C, more typically in excess of about 55° C,  
preferably in excess of about 65° C, and more preferably in  
excess of about 70° C. Stringent salt conditions will  
ordinarily be less than about 1000 mM, usually less than  
20 about 500 mM, more usually less than about 400 mM,  
typically less than about 300 mM, preferably less than  
about 200 mM, and more preferably less than about 150 mM.  
However, the combination of parameters is much more  
important than the measure of any single parameter. See,  
25 e.g., Wetmur and Davidson (1968) J. Mol. Biol. 31:349-370.

Corresponding chemokines from other mammalian species  
can be cloned and isolated by cross-species hybridization  
of closely related species. Alternatively, sequences from  
a data base may be recognized as having similarity.  
30 Homology may be relatively low between distantly related  
species, and thus hybridization of relatively closely  
related species is advisable. Alternatively, preparation  
of an antibody preparation which exhibits less species  
specificity may be useful in expression cloning approaches.  
35 PCR approaches using segments of conserved sequences will  
also be used.



## VII. Making chemokines, receptors; Mimetics

DNA which encodes each respective chemokine or fragments thereof can be obtained by chemical synthesis, screening cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples.

This DNA can be expressed in a wide variety of host cells for the synthesis of a full-length ligand or fragments which can in turn, for example, be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified molecules; and for structure/function studies. Each antigen or its fragments can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These molecules can be substantially purified to be free of protein or cellular contaminants, other than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier and/or diluent. The antigen, or portions thereof, may be expressed as fusions with other proteins.

Expression vectors are typically self-replicating DNA or RNA constructs containing the desired antigen gene or its fragments, usually operably linked to suitable genetic control elements that are recognized in a suitable host cell. These control elements are capable of effecting expression within a suitable host. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription, transcription enhancers to elevate the level of mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription

and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently of the host cell.

The vectors of this invention contain DNA which encode  
5 embodiments of a chemokine, receptor, or a fragment thereof, typically encoding a biologically active polypeptide. The DNA can be under the control of a viral promoter and can encode a selection marker. This invention further contemplates use of such expression vectors which  
10 are capable of expressing eukaryotic cDNA coding for each chemokine in a prokaryotic or eukaryotic host, where the vector is compatible with the host and where the eukaryotic cDNA coding for the ligand is inserted into the vector such that growth of the host containing the vector expresses the  
15 cDNA in question. Usually, expression vectors are designed for stable replication in their host cells or for amplification to greatly increase the total number of copies of the desirable gene per cell. It is not always necessary to require that an expression vector replicate in  
20 a host cell, e.g., it is possible to effect transient expression of the ligand or its fragments in various hosts using vectors that do not contain a replication origin that is recognized by the host cell. It is also possible to use vectors that cause integration of a chemokine gene or its  
25 fragments into the host DNA by recombination, or to integrate a promoter which controls expression of an endogenous gene.

Vectors, as used herein, comprise plasmids, viruses, bacteriophage, integratable DNA fragments, and other  
30 vehicles which enable the integration of DNA fragments into the genome of the host. Expression vectors are specialized vectors which contain genetic control elements that effect expression of operably linked genes. Plasmids are the most commonly used form of vector but all other forms of vectors  
35 which serve an equivalent function and which are, or become, known in the art are suitable for use herein. See, e.g., Pouwels, et al. (1985 and Supplements) Cloning

Vectors: A Laboratory Manual, Elsevier, N.Y., and Rodriguez, et al. (1988) (eds.) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth, Boston, MA.

5 Transformed cells include cells, preferably mammalian, that have been transformed or transfected with a chemokine gene containing vector constructed using recombinant DNA techniques. Transformed host cells usually express the ligand or its fragments, but for purposes of cloning,  
10 amplifying, and manipulating its DNA, do not need to express the protein. This invention further contemplates culturing transformed cells in a nutrient medium, thus permitting the protein to accumulate in the culture. The protein can be recovered, either from the culture or from  
15 the culture medium.

For purposes of this invention, DNA sequences are operably linked when they are functionally related to each other. For example, DNA for a presequence or secretory leader is operably linked to a polypeptide if it is  
20 expressed as a preprotein or participates in directing the polypeptide to the cell membrane or in secretion of the polypeptide. A promoter is operably linked to a coding sequence if it controls the transcription of the polypeptide; a ribosome binding site is operably linked to  
25 a coding sequence if it is positioned to permit translation. Usually, operably linked means contiguous and in reading frame, however, certain genetic elements such as repressor genes are not contiguously linked but still bind to operator sequences that in turn control expression.

30 Suitable host cells include prokaryotes, lower eukaryotes, and higher eukaryotes. Prokaryotes include both gram negative and gram positive organisms, e.g., *E. coli* and *B. subtilis*. Lower eukaryotes include yeasts, e.g., *S. cerevisiae* and *Pichia*, and species of the genus  
35 *Dictyostelium*. Higher eukaryotes include established tissue culture cell lines from animal cells, both of

non-mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host-vector systems include a wide variety of vectors for many different species. As used herein, E. coli and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A representative vector for amplifying DNA is pBR322 or many of its derivatives. Vectors that can be used to express these chemokines or their fragments include, but are not limited to, such vectors as those containing the lac promoter (pUC-series); trp promoter (pBR322-trp); Ipp promoter (the pIN-series); lambda-pP or pR promoters (pOTS); or hybrid promoters such as ptac (pDR540). See Brosius, et al. (1988) "Expression Vectors Employing Lambda-, trp-, lac-, and Ipp-derived Promoters", in Rodriguez and Denhardt (eds.) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth, Boston, Chapter 10, pp. 205-236.

Lower eukaryotes, e.g., yeasts and Dictyostelium, may be transformed with chemokine sequence containing vectors. For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, Saccharomyces cerevisiae. It will be used to generically represent lower eukaryotes although a number of other strains and species are also available. Yeast vectors typically consist of a replication origin (unless of the integrating type), a selection gene, a promoter, DNA encoding the desired protein or its fragments, and sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors for yeast include such constitutive promoters as 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionine promoter. Suitable vectors include derivatives of the following types: self-replicating low copy number (such as the YRp-series), self-replicating high copy number (such as the YEpl-series); integrating types

(such as the YIp-series), or mini-chromosomes (such as the YCp-series).

Higher eukaryotic tissue culture cells are the preferred host cells for expression of the functionally active chemokine protein. In principle, most any higher eukaryotic tissue culture cell line is workable, e.g., insect baculovirus expression systems, whether from an invertebrate or vertebrate source. However, mammalian cells are preferred, in that the processing, both cotranslationally and posttranslationally. Transformation or transfection and propagation of such cells has become a routine procedure. Examples of useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines. Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (if genomic DNA is used), a polyadenylation site, and a transcription termination site. These vectors also usually contain a selection gene or amplification gene. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as from adenovirus, SV40, parvoviruses, vaccinia virus, or cytomegalovirus. Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) Mol. Cell Biol. 5:1136-1142; pMC1neo Poly-A, see Thomas, et al. (1987) Cell 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610.

It will often be desired to express a chemokine polypeptide in a system which provides a specific or defined glycosylation pattern. In this case, the usual pattern will be that provided naturally by the expression system. However, the pattern will be modifiable by exposing the polypeptide, e.g., an unglycosylated form, to appropriate glycosylating proteins introduced into a heterologous expression system. For example, a chemokine gene may be co-transformed with one or more genes encoding

mammalian or other glycosylating enzymes. Using this approach, certain mammalian glycosylation patterns will be achievable or approximated in prokaryote or other cells.

5 A chemokine, or a fragment thereof, may be engineered to be phosphatidyl inositol (PI) linked to a cell membrane, but can be removed from membranes by treatment with a phosphatidyl inositol cleaving enzyme, e.g., phosphatidyl inositol phospholipase-C. This releases the antigen in a biologically active form, and allows purification by  
10 standard procedures of protein chemistry. See, e.g., Low (1989) Biochim. Biophys. Acta 988:427-454; Tse, et al. (1985) Science 230:1003-1008; and Brunner, et al. (1991) J. Cell Biol. 114:1275-1283.

Now that these chemokines have been characterized,  
15 fragments or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) The Practice of Peptide Synthesis, Springer-Verlag, New York;  
20 and Bodanszky (1984) The Principles of Peptide Synthesis, Springer-Verlag, New York. For example, an azide process, an acid chloride process, an acid anhydride process, a mixed anhydride process, an active ester process (for  
25 example, p-nitrophenyl ester, N-hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimidazole process, an oxidative-reductive process, or a dicyclohexylcarbodiimide (DCCD)/additive process can be used. Solid phase and solution phase syntheses are both applicable to the  
30 foregoing processes.

These chemokines, fragments, or derivatives are suitably prepared in accordance with the above processes as typically employed in peptide synthesis, generally either by a so-called stepwise process which comprises condensing  
35 an amino acid to the terminal amino acid, one by one in sequence, or by coupling peptide fragments to the terminal amino acid. Amino groups that are not being used in the

coupling reaction are typically protected to prevent coupling at an incorrect location.

If a solid phase synthesis is adopted, the C-terminal amino acid is typically bound to an insoluble carrier, or support through its carboxyl group. The insoluble carrier is not particularly limited as long as it has a binding capability to a reactive carboxyl group. Examples of such insoluble carriers include halomethyl resins, such as chloromethyl resin or bromomethyl resin, hydroxymethyl resins, phenol resins, tert-alkyloxycarbonyl-hydrazidated resins, and the like.

An amino group-protected amino acid is bound in sequence through condensation of its activated carboxyl group and the reactive amino group of the previously formed peptide or chain, to synthesize the peptide step by step. After synthesizing the complete sequence, the peptide is split off from the insoluble carrier to produce the peptide. This solid-phase approach is generally described by Merrifield, et al. (1963) in J. Am. Chem. Soc. 85:2149-2156.

The prepared ligand and fragments thereof can be isolated and purified from the reaction mixture by means of peptide separation, e.g., by extraction, precipitation, electrophoresis and various forms of chromatography, and the like. The various chemokines of this invention can be obtained in varying degrees of purity depending upon its desired use. Purification can be accomplished by use of the protein purification techniques disclosed herein or by the use of the antibodies herein described, e.g., in immunoabsorbant affinity chromatography. This immunoabsorbant affinity chromatography is carried out by first linking the antibodies to a solid support and then contacting the linked antibodies with solubilized lysates of appropriate source cells, lysates of other cells expressing the ligand, or lysates or supernatants of cells producing the desired chemokine as a result of DNA techniques, see below.

## VIII. Uses

The present invention provides reagents which will find use in diagnostic applications as described elsewhere herein, e.g., in the general description for developmental abnormalities, or below in the description of kits for diagnosis.

This invention also provides reagents with significant therapeutic value. These chemokines (naturally occurring or recombinant), fragments thereof and antibodies thereto, along with compounds identified as having binding affinity to them, should be useful in the treatment of conditions associated with abnormal physiology or development, including inflammatory conditions, including asthma. In particular, modulation of trafficking of leukocytes is one likely biological activity, but a wider tissue distribution might suggest broader biological activity, including, e.g., antiviral effects. Abnormal proliferation, regeneration, degeneration, and atrophy may be modulated by appropriate therapeutic treatment using the compositions provided herein. For example, a disease or disorder associated with abnormal expression or abnormal signaling by a chemokine should be a likely target for an agonist or antagonist of the ligand.

Various abnormal physiological or developmental conditions are known in cell types shown to possess the chemokine mRNAs by Northern blot analysis. See Berkow (ed.) The Merck Manual of Diagnosis and Therapy, Merck & Co., Rahway, N.J.; and Thorn, et al. Harrison's Principles of Internal Medicine, McGraw-Hill, N.Y. Developmental or functional abnormalities, e.g., of the immune system, cause significant medical abnormalities and conditions which may be susceptible to prevention or treatment using compositions provided herein.

Chemokine antibodies, including recombinant forms, can be purified and then administered to a patient. These reagents can be combined for therapeutic use with



additional active or inert ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, e.g., immunogenic adjuvants, along with physiologically innocuous stabilizers and excipients.

- 5 These combinations can be sterile filtered and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates use of antibodies or binding fragments thereof, including forms which are not complement binding.
- 10 Moreover, modifications to the antibody molecules or antigen binding fragments thereof, may be adopted which affect the pharmacokinetics or pharmacodynamics of the therapeutic entity.

- Drug screening using antibodies or receptor or
- 15 fragments thereof can be performed to identify compounds having binding affinity to each chemokine or receptor, including isolation of associated components. Subsequent biological assays can then be utilized to determine if the compound has intrinsic stimulating activity and is
- 20 therefore a blocker or antagonist in that it blocks the activity of the ligand. Likewise, a compound having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of a chemokine. This invention further contemplates the
- 25 therapeutic use of antibodies to these chemokines as antagonists. This approach should be particularly useful with other chemokine species variants.

- The quantities of reagents necessary for effective therapy will depend upon many different factors, including
- 30 means of administration, target site, physiological state of the patient, and other medicants administered. Thus, treatment dosages should be titrated to optimize safety and efficacy in various populations, including racial subgroups, age, gender, etc. Typically, dosages used in
- 35 vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders

will provide further predictive indication of human dosage. Various considerations are described, e.g., in Gilman, et al. (eds.) (1990) Goodman and Gilman's: The

Pharmacological Bases of Therapeutics, 8th Ed., Pergamon

5 Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.. Methods for administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others.

10 Pharmaceutically acceptable carriers typically include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, New Jersey. Dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about  
15 10  $\mu$ M concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or a slow release apparatus will often be utilized for continuous  
20 administration.

A chemokine, fragments thereof, or antibodies to it or its fragments, antagonists, and agonists, may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable  
25 to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in any conventional dosage formulation. While it is possible for the active ingredient to be administered alone, it is often preferable  
30 to present it as a pharmaceutical formulation.

Formulations typically comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier should be both pharmaceutically and physiologically acceptable in the  
35 sense of being compatible with the other ingredients and not injurious to the patient. Carriers may improve storage life, stability, etc. Formulations include those suitable

for oral, rectal, nasal, or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds.) (1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack Publishing Co., Easton, Penn.; Avis, et al. (eds.) (1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, New York; Lieberman, et al. (eds.) (1990) Pharmaceutical Dosage Forms: Tablets Dekker, New York; and Lieberman, et al. (eds.) (1990) Pharmaceutical Dosage Forms: Disperse Systems Dekker, New York. The therapy of this invention may be combined with or used in association with other therapeutic agents.

Both the naturally occurring and the recombinant forms of the chemokines of this invention are particularly useful in kits and assay methods which are capable of screening compounds for binding activity to the proteins. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds in a short period. See, e.g., Fodor, et al. (1991) Science 251:767-773, which describes means for testing of binding affinity by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays can be greatly facilitated by the availability of large amounts of purified, soluble chemokine as provided by this invention.

For example, antagonists can normally be found once the ligand has been structurally defined. Testing of potential ligand analogs is now possible upon the development of highly automated assay methods using physiologically responsive cells. In particular, new agonists and antagonists will be discovered by using screening techniques described herein.

Viabile cells could also be used to screen for the effects of drugs on respective chemokine mediated functions, e.g., second messenger levels, i.e.,  $\text{Ca}^{++}$ ; inositol phosphate pool changes (see, e.g., Berridge (1993) Nature 361:315-325 or Billah and Anthes (1990) Biochem. J. 269:281-291); cellular morphology modification responses; phosphoinositide lipid turnover; an antiviral response. and others. Some detection methods allow for elimination of a separation step, e.g., a proximity sensitive detection system. Calcium sensitive dyes will be useful for detecting  $\text{Ca}^{++}$  levels, with a fluorimeter or a fluorescence cell sorting apparatus.

Rational drug design may also be based upon structural studies of the molecular shapes of the chemokines and other effectors or analogs. Effectors may be other proteins which mediate other functions in response to ligand binding, or other proteins which normally interact with the receptor. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) Protein Crystallography, Academic Press, New York.

Purified chemokine can be coated directly onto plates for use in the aforementioned drug screening techniques. However, non-neutralizing antibodies to these ligands can be used as capture antibodies to immobilize the respective ligand on the solid phase.

Similar concepts also apply to the chemokine receptor embodiments of the invention.

#### IX. Kits

This invention also contemplates use of chemokine proteins, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for

detecting the presence of ligand, antibodies, or chemokine receptors. Typically the kit will have a compartment containing either a defined chemokine peptide or gene segment or a reagent which recognizes one or the other, e.g., antibodies.

A kit for determining the binding affinity of a test compound to a chemokine would typically comprise a test compound; a labeled compound, for example an antibody having known binding affinity for the ligand; a source of chemokine (naturally occurring or recombinant); and a means for separating bound from free labeled compound, such as a solid phase for immobilizing the ligand. Once compounds are screened, those having suitable binding affinity to the ligand can be evaluated in suitable biological assays, as are well known in the art, to determine whether they act as agonists or antagonists to the receptor. The availability of recombinant chemokine polypeptides also provide well defined standards for calibrating such assays or as positive control samples.

A preferred kit for determining the concentration of, for example, a chemokine in a sample would typically comprise a labeled compound, e.g., antibody, having known binding affinity for the ligand, a source of ligand (naturally occurring or recombinant) and a means for separating the bound from free labeled compound, for example, a solid phase for immobilizing the chemokine. Compartments containing reagents, and instructions for use or disposal, will normally be provided.

Antibodies, including antigen binding fragments, specific for the chemokine or ligand fragments are useful in diagnostic applications to detect the presence of elevated levels of chemokine and/or its fragments. Such diagnostic assays can employ lysates, live cells, fixed cells, immunofluorescence, cell cultures, body fluids, and further can involve the detection of antigens related to the ligand in serum, or the like. Diagnostic assays may be homogeneous (without a separation step between free reagent

and antigen-ligand complex) or heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied  
5 immunoassay technique (EMIT), substrate-labeled fluorescent immunoassay (SLFIA), and the like. For example, unlabeled antibodies can be employed by using a second antibody which is labeled and which recognizes the antibody to a chemokine or to a particular fragment thereof. Similar assays have  
10 also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH.

Anti-idiotypic antibodies may have similar uses to diagnose presence of antibodies against a chemokine, as  
15 such may be diagnostic of various abnormal states. For example, overproduction of a chemokine may result in production of various immunological reactions which may be diagnostic of abnormal physiological states, particularly in various inflammatory or asthma conditions.

20 Frequently, the reagents for diagnostic assays are supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody or labeled chemokine is  
25 provided. This is usually in conjunction with other additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the  
30 contents after use. Typically the kit has compartments for each useful reagent. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium providing appropriate concentrations of reagents for performing the assay.

35 The aforementioned constituents of the drug screening and the diagnostic assays may be used without modification or may be modified in a variety of ways. For example,

labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, the ligand, test compound, chemokine, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as  $^{125}\text{I}$ , enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating bound from the free ligand, or alternatively bound from free test compound. The chemokine can be immobilized on various matrixes followed by washing. Suitable matrixes include plastic such as an ELISA plate, filters, and beads. Methods of immobilizing the chemokine to a matrix include, without limitation, direct adhesion to plastic, use of a capture antibody, chemical coupling, and biotin-avidin. The last step in this approach involves the precipitation of ligand/antibody complex by any of several methods including those utilizing, e.g., an organic solvent such as polyethylene glycol or a salt such as ammonium sulfate. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) Clin. Chem. 30:1457-1461, and the double antibody magnetic particle separation as described in U.S. Pat. No. 4,659,678.

Methods for linking proteins or their fragments to the various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of activated carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by reaction of a mercapto group with an

activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves  
5 use of oligonucleotide or polynucleotide sequences taken from the sequence of a chemokine. These sequences can be used as probes for detecting levels of the ligand message in samples from patients suspected of having an abnormal condition, e.g., an inflammatory or developmental problem.  
10 The preparation of both RNA and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides, usually at least  
15 about 18 nucleotides, and the polynucleotide probes may be up to several kilobases. Various labels may be employed, most commonly radionuclides, particularly  $^{32}\text{P}$ . However, other techniques may also be employed, such as using biotin modified nucleotides for introduction into a  
20 polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as radionuclides, fluorescers, enzymes, or the like. Alternatively, antibodies may be employed which can recognize specific  
25 duplexes, including DNA duplexes, RNA duplexes, DNA-RNA hybrid duplexes, or DNA-protein duplexes. The antibodies in turn may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to  
30 the duplex can be detected. The use of probes to the novel anti-sense RNA may be carried out in any conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational probing, hybrid released translation (HRT), and hybrid arrested translation (HART).  
35 This also includes amplification techniques such as polymerase chain reaction (PCR).



Diagnostic kits which also test for the qualitative or quantitative presence of other markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers. Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) Progress in Growth Factor Res. 1:89-97.

#### X. Receptor

Having isolated a ligand binding partner of a specific interaction, methods exist for isolating the counter-partner. See, Gearing, et al EMBO J. 8:3667-4676 or McMahan, et al. (1991) EMBO J. 10:2821-2832. For example, means to label a chemokine without interfering with the binding to its receptor can be determined. For example, an affinity label can be fused to either the amino- or carboxy-terminus of the ligand. An expression library can be screened for specific binding of chemokine, e.g., by cell sorting, or other screening to detect subpopulations which express such a binding component. See, e.g., Ho, et al. (1993) Proc. Nat'l Acad. Sci. 90:11267-11271. Alternatively, a panning method may be used. See, e.g., Seed and Aruffo (1987) Proc. Nat'l. Acad. Sci. 84:3365-3369.

Protein cross-linking techniques with label can be applied to a isolate binding partners of a chemokine. This would allow identification of protein which specifically interacts with a chemokine, e.g., in a ligand-receptor like manner.

In various embodiments, new receptors designated DC CR and M/DC CR were isolated. The sequences of the human constructs and product are provided in Tables 4 and 5. Similar means for making variants and fragments, at the nucleotide level or at the protein level, and making antibodies will be available as described above, directed primarily to the chemokine embodiments. Many similar or

related uses to the ligands will be applied to the  
receptors, as specific binding reagents. In particular,  
methods will be applied to screening for specific ligands  
for each receptor. Many uses, including kits, will also be  
5 available through analogous techniques.

The broad scope of this invention is best understood  
with reference to the following examples, which are not  
intended to limit the invention to specific embodiments.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
220

EXAMPLES

## I. General Methods

- 5        Some of the standard methods are described or  
referenced, e.g., in Maniatis, et al. (1982) Molecular  
Cloning, A Laboratory Manual, Cold Spring Harbor  
Laboratory, Cold Spring Harbor Press; Sambrook, et al.  
(1989) Molecular Cloning: A Laboratory Manual, (2d ed.),  
10 vols 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene  
Publishing Associates, Brooklyn, NY; or Ausubel, et al.  
(1987 and Supplements) Current Protocols in Molecular  
Biology, Greene/Wiley, New York; Innis, et al. (eds.) (1990)  
PCR Protocols: A Guide to Methods and Applications Academic  
15 Press, N.Y. Methods for protein purification include such  
methods as ammonium sulfate precipitation, column  
chromatography, electrophoresis, centrifugation,  
crystallization, and others. See, e.g., Ausubel, et al.  
(1987 and periodic supplements); Deutscher (1990) "Guide to  
20 Protein Purification" in Methods in Enzymology, vol. 182,  
and other volumes in this series; and manufacturer's  
literature on use of protein purification products, e.g.,  
Pharmacia, Piscataway, N.J., or Bio-Rad, Richmond, CA.  
Combination with recombinant techniques allow fusion to  
25 appropriate segments, e.g., to a FLAG sequence or an  
equivalent which can be fused via a protease-removable  
sequence. See, e.g., Hochuli (1989) Chemische Industrie  
12:69-70; Hochuli (1990) "Purification of Recombinant  
Proteins with Metal Chelate Absorbent" in Setlow (ed.)  
30 Genetic Engineering, Principle and Methods 12:87-98, Plenum  
Press, N.Y.; and Crowe, et al. (1992) OIAexpress: The High  
Level Expression & Protein Purification System QUIAGEN,  
Inc., Chatsworth, CA.
- FACS analyses are described in Melamed, et al. (1990)  
35 Flow Cytometry and Sorting Wiley-Liss, Inc., New York, NY;  
Shapiro (1988) Practical Flow Cytometry Liss, New York, NY;

and Robinson, et al. (1993) Handbook of Flow Cytometry Methods Wiley-Liss, New York, NY.

## II. Isolation and characterization of chemokine cDNAs

### 5 A. TECK

The TECK was isolated from a cDNA library made from thymus cells from a RAG-1 "knockout" mouse. See, Mombaerts, et al. (1992) Cell 68:869-877. Individual cDNA clones were sequenced using standard methods, e.g., the Taq  
10 DyeDeoxy Terminator Cycle Sequencing kit (Applied Biosystems, Foster City, CA), and the TECK sequence was identified and further characterized. Computer analyses with other C-C chemokine family members revealed  
15 significant homology at the amino acid levels with other chemokines. The nucleotide sequence for mouse is provided in Table 1, encoding a polypeptide of about 144 amino acids. The signal sequence should run from 1 (met) to about 23 (ala), and removal of the signal sequence should provide one natural mature sequence beginning at 24 (gln).  
20 Additional processing may occur in a physiological system.

The sequence is notable in having a longer carboxy-terminal tail than most other CC chemokines. TECK exhibits one glycosylation site, and several AAMP, PKC, and CK2 phosphorylation sites.

### 25 B. MIP-3 $\alpha$

The MIP-3 $\alpha$  was isolated from a cDNA library made from human monocytes activated with LPS and IFN- $\gamma$ . Individual cDNA clones were sequenced using standard methods, and the MIP-3 $\alpha$  sequence was identified and further characterized.  
30 The nucleotide sequence is provided in Table 2, encoding a polypeptide of at least about 89 amino acids. The signal sequence should run from about 1 (met) to 21 (cys), and removal of the signal sequence should provide one natural sequence beginning with gly. Additional processing may  
35 occur in a physiological system.

C. MIP-3 $\beta$ 

The MIP-3 $\beta$  was isolated from a cDNA library made from human fetal lung cells. Individual cDNA clones are sequenced using standard methods, and the MIP-3 $\alpha$  sequence was identified and further characterized. The nucleotide sequence is provided in Table 3, encoding a polypeptide of about 98 amino acids. The signal sequence should run from about 1 (met) to about 21 (ser), and removal of the signal sequence should provide one mature natural sequence beginning from gly. Additional processing may occur in a physiological system.

This chemokine has been paired with a receptor designated Ebil. See Yoshida, et al. (1997) J. Biol. Chem. 13803-13809.

## 15 D. Dendritic Cell Receptor for chemokine; DC CR

The DC CR was isolated from RNA made from dendritic cells isolated from CD34<sup>+</sup> cord blood cells, isolated by standard procedure. It was also isolated from eosinophils using degenerate PCR primers of the TM2 and TM7 segments, which are often conserved among chemokine receptors. These eosinophils were isolated by taking PBLs, depletion of red blood cells by lysis, and negative selection of CD16 to remove neutrophils.

Sequencing of the PCR fragments indicated a potential novel receptor, and the fragment was used to isolate a full length clone by hybridization. Clone isolates were sequenced using standard methods, and the DC CR sequence was identified and further characterized. The nucleotide sequence is provided in Table 4, encoding a polypeptide of about 365 amino acids. The transmembrane segments, determined by homology to the IL-8 B receptor, are about: TM1 from 39 (leu) to 64 (phe); TM2 from 76 (leu) to 96 (ser); TM3 from 111 (leu) to 132 (met); TM4 from 151 (thr) to 176 (phe); TM5 from 207 (gly) to 229 (val); TM6 from 246 (val) to 270 (ala); and TM7 from 291 (val) to 319 (leu). The amino terminal segment is probably an extracellular

segment, and the others would be between TM2 and TM3; and TM4 and TM5; and TM6 and TM7. The intracellular segments should then run between TM1 and TM2; TM3 and TM4, TM5 and TM6, and the carboxy terminus from the end of TM7.

5 Additional processing may occur in a physiological system.

The implication of chemokine receptors in retroviral infection suggest that the receptor may be critical for infection. Antibodies which block infection may be routinely screened, and developed for therapeutic uses.

10 E. Monocyte/Dendritic Cell Receptor for chemokine;  
M/DC CR

The M/DC CR was isolated from a cDNA library made from human monocyte cells cultured for 2.5 to 4 h in medium containing IFN- $\gamma$  (10 ng/ml), LPS (1  $\mu$ g/ml), anti-IL-4  
15 monoclonal antibody (5  $\mu$ g/ml), and anti-IL-10 monoclonal antibody (5  $\mu$ g/ml). Individual cDNA clones were sequenced using standard methods, and the M/DC CR sequence was identified and further characterized. The nucleotide sequence is provided in Table 5, encoding a polypeptide of  
20 about 356 amino acids. The transmembrane segments, should be about as follows: TM1 from 52 (leu) to 76 (val); TM2 from 86 (asn) to 107 (ala); TM3 from 117 (ile) to 138 (val); TM4 from 157 (val) to 182 (tyr); TM5 from 211 (phe) to 233 (val); TM6 from 251 (leu) to 275 (phe); and TM7 from  
25 296 (ile) to 315 (leu). As for the DC CR, the amino terminal segment is probably an extracellular segment, and the others would be between TM2 and TM3; and TM4 and TM5; and TM6 and TM7. The intracellular segments should then run between TM1 and TM2; TM3 and TM4, TM5 and TM6, and the  
30 carboxy terminus from the end of TM7.

### III. Preparation of antibodies

Many standard methods are available for preparation of antibodies. For example, synthetic peptides may be  
35 prepared to be used as antigen, administered to an appropriate animal, and either polyclonal or monoclonal antibodies prepared. Short peptides, e.g., less than about

10 amino acids may be repeated, while longer peptides may be used alone or conjugated to a carrier. For example, with the M/DC CR, animals were immunized with peptides corresponding to amino acid sequences from 18-44 (starting with LAP and ending with KYD; a fragment towards the amino terminus) and from 183-204 (starting with KPQ and ending with PAD; corresponding to an extracellular loop), see SEQ ID NO: 13. Highest specificity will result when the polypeptides are selected from portions which are most unique, e.g., not from conserved sequence regions. The animals may be used to collect antiserum, or may be used to generate monoclonal antibodies.

Antiserum was determined useful for ELISA, and will be evaluated for utility as immunoprecipitation or Western blot analysis. Monoclonal antibodies will also be evaluated for those same uses.

The antibodies provided will be useful as immunoaffinity reagents, as detection reagents, for immunohistochemistry, and as therapeutic reagents.

#### IV. Assays for chemotactic activity of chemokines.

Chemokine proteins are produced, e.g., in COS cells transfected with a plasmid carrying the chemokine cDNA by electroporation. See, Hara, et al. (1992) EMBO J. 10:1875-1884. Physical analytical methods may be applied, e.g., CD analysis, to compare tertiary structure to other chemokines to evaluate whether the protein has likely folded into an active conformation. After transfection, a culture supernatant is collected and subjected to bioassays. A mock control, e.g., a plasmid carrying the luciferase cDNA, is used. See, de Wet, et al. (1987) Mol. Cell. Biol. 7:725-757. A positive control, e.g., recombinant murine MIP-1 $\alpha$  from R&D Systems (Minneapolis, MN), is typically used. Likewise, antibodies may be used to block the biological activities, e.g., as a control.

Lymphocyte migration assays are performed as previously described, e.g., in Bacon, et al. (1988) Br. J.

Pharmacol. 95:966-974. Murine Th2 T cell clones, CDC-25 (see Tony, et al. (1985) J. Exp. Med. 161:223-241) and HDK-1 (see Cherwinski, et al. (1987) J. Exp. Med. 166:1229-1244), made available from R. Coffman and A.

- 5 O'Garra (DNAX, Palo Alto, CA), respectively, are used as controls.

Ca<sup>2+</sup> flux upon chemokine stimulation is measured according to the published procedure described in Bacon, et al. (1995) J. Immunol. 154:3654-3666.

- 10 Maximal numbers of migrating cells in response to MIP-1 $\alpha$  typically occur at a concentration of 10<sup>-8</sup> M, in agreement with original reports for CD4<sup>+</sup> populations of human T cells. See Schall (1993) J. Exp. Med. 177:1821-1826. A dose-response curve is determined,  
15 preferably giving a characteristic bell shaped dose-response curve.

- After stimulation with C-C chemokines, lymphocytes generally show a measurable intracellular Ca<sup>2+</sup> flux. MIP-1 $\alpha$  is capable of inducing immediate transients of calcium  
20 mobilization. Typically, the levels of chemokine used in these assays will be comparable to those used for the chemotaxis assays (1/1000 dilution of conditioned supernatants).

- Retroviral infection assays have also been described,  
25 and recent description of certain chemokine receptors in retroviral infection processes may indicate that similar roles may apply to the DC CR and/or M/DC CR. See, e.g., Balter (1996) Science 272:1740 (describing evidence for chemokine receptors as coreceptors for HIV); and Deng, et  
30 al. (1996) Nature 381:661-666.

#### V. Expression analysis of chemokine/receptor genes

- RNA blot and hybridization are performed according to the standard method in Maniatis, et al. (1982) Molecular  
35 Cloning: A Laboratory Manual Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. An appropriate fragment of a cDNA fragment is selected for use as a probe. To verify



the amount of RNA loaded in each lane, the substrate membrane is reprobbed with a control cDNA, e.g., glyceraldehyde 3-phosphate dehydrogenase (G3PDH) cDNA (Clontech, Palo Alto CA).

5        Analysis of mRNA from the appropriate cell source using the probe will determine the natural size of message. It will also indicate whether different sized messages exist. The messages will be subject to analysis after isolation, e.g., by PCR or hybridization techniques.

10       Northern blot analysis may be performed on many different mRNA sources. However, in certain cases, cDNA libraries may be used to evaluate sources which are difficult to prepare. A "reverse Northern" uses cDNA inserts removed from vector, but multiplicity of bands may  
15       reflect either different sized messages, or may be artifact due to incomplete reverse transcription in the preparation of the cDNA library. In such instances, verification may be appropriate by standard Northern analysis.

20       Similarly, Southern blots may be used to evaluate species distribution of a gene. The stringency of washes of the blot will also provide information as to the extent of homology of various species counterparts.

Tissue distribution, and cell distribution, may be evaluated by immunohistochemistry using antibodies.  
25       Alternatively, in situ nucleic acid hybridization may also be used in such analysis.

#### A. TECK

The TECK was isolated from a RAG-1 "knockout" mouse. This animal is characterized by a great predominance of  
30       pro-T or pre-T cells, lacking more mature T cells after the block point of T cell receptor rearrangement. This suggests a role in very early T cell development, likely expressed by pro-T or pre-T cells, thymic stromal cells, and possibly macrophages, epithelial, and dendritic cells.  
35       This comports with the observation that tissue distribution studies have not detected significant expression in other organs or tissues. See also, Table 7.

Variable	Mean	SD	Min	Max	Skewness	Kurtosis	Normality
Age	38.5	12.5	25	65	0.1	-0.2	0.95
Gender	1.2	0.4	1	2	0.0	0.0	0.99
Marital Status	1.8	0.8	1	3	0.2	0.1	0.98
Education	15.2	2.1	10	20	0.3	0.2	0.97
Income	1200	300	500	2000	0.4	0.3	0.96
Occupation	1.5	0.5	1	3	0.1	0.0	0.99
Health Status	1.5	0.5	1	3	0.1	0.0	0.99
Stress Level	2.5	0.8	1	4	0.2	0.1	0.98
Life Satisfaction	3.5	0.9	1	5	0.3	0.2	0.97
Resilience	2.8	0.7	1	4	0.2	0.1	0.98
Emotional Stability	3.2	0.8	1	5	0.3	0.2	0.97
Relationship Quality	3.8	0.9	1	5	0.3	0.2	0.97
Parenting Style	2.2	0.6	1	4	0.2	0.1	0.98
Child Development	2.5	0.7	1	4	0.2	0.1	0.98
Family Cohesion	3.0	0.8	1	5	0.3	0.2	0.97
Parental Involvement	3.5	0.9	1	5	0.3	0.2	0.97
Child Behavior	2.0	0.6	1	4	0.2	0.1	0.98
Family Functioning	3.2	0.8	1	5	0.3	0.2	0.97
Parental Stress	2.5	0.7	1	4	0.2	0.1	0.98
Child Health	3.0	0.8	1	5	0.3	0.2	0.97
Family Support	3.5	0.9	1	5	0.3	0.2	0.97
Parental Satisfaction	3.8	0.9	1	5	0.3	0.2	0.97
Child Well-being	3.0	0.8	1	5	0.3	0.2	0.97
Family Resilience	3.2	0.8	1	5	0.3	0.2	0.97
Parental Efficacy	3.5	0.9	1	5	0.3	0.2	0.97
Child Social Skills	2.5	0.7	1	4	0.2	0.1	0.98
Family Communication	3.0	0.8	1	5	0.3	0.2	0.97
Parental Involvement	3.5	0.9	1	5	0.3	0.2	0.97
Child Development	2.5	0.7	1	4	0.2	0.1	0.98
Family Cohesion	3.0	0.8	1	5	0.3	0.2	0.97
Parental Involvement	3.5	0.9	1	5	0.3	0.2	0.97
Child Behavior	2.0	0.6	1	4	0.2	0.1	0.98
Family Functioning	3.2	0.8	1	5	0.3	0.2	0.97
Parental Stress	2.5	0.7	1	4	0.2	0.1	0.98
Child Health	3.0	0.8	1	5	0.3	0.2	0.97
Family Support	3.5	0.9	1	5	0.3	0.2	0.97
Parental Satisfaction	3.8	0.9	1	5	0.3	0.2	0.97
Child Well-being	3.0	0.8	1	5	0.3	0.2	0.97
Family Resilience	3.2	0.8	1	5	0.3	0.2	0.97
Parental Efficacy	3.5	0.9	1	5	0.3	0.2	0.97
Child Social Skills	2.5	0.7	1	4	0.2	0.1	0.98
Family Communication	3.0	0.8	1	5	0.3	0.2	0.97
Parental Involvement	3.5	0.9	1	5	0.3	0.2	0.97
Child Development	2.5	0.7	1	4	0.2	0.1	0.98
Family Cohesion	3.0	0.8	1	5	0.3	0.2	0.97
Parental Involvement	3.5	0.9	1	5	0.3	0.2	0.97
Child Behavior	2.0	0.6	1	4	0.2	0.1	0.98
Family Functioning	3.2	0.8	1	5	0.3	0.2	0.97
Parental Stress	2.5	0.7	1	4	0.2	0.1	0.98
Child Health	3.0	0.8	1	5	0.3	0.2	0.97
Family Support	3.5	0.9	1	5	0.3	0.2	0.97
Parental Satisfaction	3.8	0.9	1	5	0.3	0.2	0.97
Child Well-being	3.0	0.8	1	5	0.3	0.2	

Table 7: mTECK mRNA expression in tissues and cells

cDNA libraries			northern blot		
cell type or tissue	neg	pos	cell type or tissue	neg	pos
Th2 CD4+ T cells	X		heart	X	
Th1 CD4+ T cells	X		brain	X	
Lung	X		spleen	X	
L cells	X		lung	X	
RAG-1 KO lung	X		liver	X	
RAG-1 KO heart	X		skeletal muscle	X	
RAG-1 KO brain		X (+)	kidney	X	
RAG-1 KO spleen	X		testis	X	
RAG-1 KO kidney	X		thymus		X (+++)
RAG-1 KO testis		X (+)	small intestine		X (++)
RAG-1 KO thymus		X (+++)	CD4+8- thymocytes R/A	X	
RAG-1 KO liver		X (+)	CD4-8+ thymocytes R/A	X	
CD4-8- thymocytes	X		CD4-8- thymocytes R/A	X	
A20-J B-cell lymphoma	X		B220+ splenocytes R/A	X	
BW CD4-8-3- hybridoma	X		Thy-1+ splenocytes R/A	X	
pro-T cells		X (+)	1G18LA macrophages R/A	X	
pre-T cells	X		primary thymic stroma R/A	X	
30-R bone marrow stroma	X		3D.1 thymic epithelial R/A	X	
D10 T-cell hybridoma	X		MTSC-C thymic epithelial	X	
CTLL T-cell clone	X		30.R bone marrow stroma	X	
peritoneal macrophages	X				
splenic dendritic cells	X				

Analysis of mTECK mRNA was carried out as described. + to +++ indicates the relative intensity of the signal. R/A: resting or activated.

Species analysis indicated positive signals by hybridization in human, rat, and hamster DNA. Tissue distribution analysis suggests that the gene is expressed in human small intestine, which also is a tissue which supports T cell differentiation.

The combination of the structure and distribution of this chemokine suggests a role in T cell development, which normally occurs in the thymus.

#### B. MIP-3 $\alpha$

The MIP-3 $\alpha$  was identified from a cDNA library made from human monocytes activated with LPS and IFN- $\gamma$ , in the presence of anti-IL-10. See, Rossi, et al. (1997) J. Immunology 158:1033-1036, which was published after priority dates of this filing. Message of the chemokine has also been detected in pancreatic islet cells, fetal lung, and hepatic HEPG2 cells, suggesting a physiological role in inflammation or medical conditions in such organs/tissues.

The gene is expressed in HL-60 (promyelocytic leukemia); S3 (HeLa cell); K562 (chronic myelogenous leukemia); MOLT-4 (lymphoblastic leukemia); Raji (Burkitt's lymphoma); SW480 (colorectal adenocarcinoma); A549 (lung carcinoma); and G361 (melanoma) cell lines, as determined by probing on a tissue blot from CLONTECH. Tissue expression gave a positive signal in lymph node, appendix, peripheral blood lymphocytes, fetal liver, and fetal lung, suggesting a physiological role in inflammation or medical conditions in such organs/tissues; but no detectable signal in spleen, bone marrow, brain, and kidney.

The main transcript appears to be about 1.2 kb, with two additional transcript sizes in fetal lung RNA. Among the various tissues, transcript sizes of 1.8, 2.7, and 4.2 kb were detected.

Positive signals were also detected in the following cDNA libraries: dendritic cells activated with LPS, but not when activated with GM-CSF and IL-4; monocytes treated

with LPS, IFN- $\gamma$ , and anti-IL-10, but not when treated with LPS, IFN- $\gamma$ , and IL-10; and activated PBMC.

These expression data implicate this chemokine in inflammatory responses upon cell activation. The lymph nodes, appendix, and PBL are sites where inflammatory processes take place. The MIP-3 $\alpha$  may exert its effects on monocytes and cells involved in inflammatory events. Other structural features implicate this chemokine in eosinophil and lung physiology, e.g., asthma indications. Thus, an antagonist of the chemokine, e.g., an antibody, may be important for treatment of asthmatic conditions. Also, IL-10 appears to inhibit MIP-3 $\alpha$  expression.

The human MIP-3 $\alpha$  is a ligand for the DC CR. Thus, a positive control exists for the the Ca<sup>++</sup> flux assay for that receptor. This allows for the further screening of agonist ligands for the DC CR. Moreover, the DC CR was isolated from eosinophil cDNA, and observations have been made that eosinophils migrate to MIP-3 $\alpha$  in vitro. These suggest that the MIP-3 $\alpha$  interaction with the DC CR is important in recruitment of eosinophils, as occurs with the eotaxin ligand and the CCR3. As such, antagonists of the MIP-3 $\alpha$  interaction with the DC CR will likely be useful in inhibiting eosinophilia, particularly in the lung, or lung inflammation. These may accompany asthmatic or other pulmonary conditions.

Antagonists to MIP-3 $\alpha$  may be made either with antibodies, or other binding compositions which inhibit receptor interaction. The antibodies may be to the ligand, MIP-3 $\alpha$  itself, or to the binding portions of the receptor, DC CR. Muteins of the chemokine may block receptor interaction, and with a positive control, chemokine muteins may be screened for variations which compete with the wild type chemokine at various concentrations. See, e.g., Kenakin (1987) Pharmacological Analysis of Drug-Receptor Interaction Raven Press, NY; Arunlakshana and Schild (1959) Br. J. Pharmacol. 14:48-58; Black (1989) Science 245:486-493; Zurawski, et al. (1986) J. Immunol. 137:3354-3360;

Zurawski and Zurawski (1988) EMBO J. 7:1061-1069; Zurawski and Zurawski (1992) EMBO J. 11:3905-3910; Imler and Zurawski (1992) J. Biol. Chem. 267:13185-13190.

### C. MIP-3 $\beta$

5       The MIP-3 $\beta$  was identified in a cDNA library made from human monocytes activated with LPS and IFN- $\gamma$ , in the presence of anti-IL-10. Its distribution in other cells and tissues has not been fully determined.

### D. Dendritic Cell Receptor for chemokine; DC CR

10       The DC CR was isolated from a cDNA library made from a dendritic cell cDNA library. It appears to be expressed in certain T cells, spleen cell subsets, NK cells, and other cell populations enriched in dendritic cells, including CD1a<sup>+</sup>, CD14<sup>+</sup>, and CD1Aa<sup>+</sup> cells. It did not give a  
15       detectable signal in TF1, Jurkat, MRC5, JY, or U937 cells.

      Being found on dendritic cells, its ligand, including the MIP-3 $\alpha$ , may be important in attracting appropriate cells for the initiation of an immune response. MIP-3 $\alpha$  has been shown to be a very potent chemoattractant for  
20       dendritic cells. Significant roles of the ligand and receptor in pulmonary physiology are suggested, both from the distribution of the receptor and ligand. The receptor may be also present in other cells important in such responses.

### 25       E. Monocyte/Dendritic Cell Receptor for chemokine; M/DC CR

      The M/DC CR was isolated from a cDNA library made from primary monocyte cells activated with LPS and IFN- $\gamma$  but subtracted with known high abundance genes from those  
30       cells. The abundance of this gene is probably less than about 1% of message from those cells.

      Tissue expression gave a positive signal in spleen, PBL, lung, placenta, and small intestine; but no detectable signal in brain, liver, kidney, and muscle. This  
35       distribution suggests a hematopoietic role.

There appears to be one main transcript, but the existence of additional or alternatively spliced messages has not been eliminated.

Positive signals were also detected in the following  
5 cDNA libraries: monocytes and dendritic cells; but signals  
were not detectable in CD8<sup>+</sup> T cells, or in either resting  
or activated splenocytes, gamma-delta T cells, NK cells, or  
B cells. Immunohistochemistry will be performed to confirm  
absence in the T cell and B cell compartments and to check  
10 in tonsil, particularly in view of location in spleen and  
placenta. The relatively restricted distribution on  
monocytes and dendritic cells leads both to its  
designation, and suggests a functional role in those cell  
types, which are important in the initiation of immune  
15 responses through their ability to process and present  
antigen to T cells.

## VI. Specific Characterization of TECK

A novel CC chemokine was identified in the thymus of  
20 mouse and human and was designated TECK as Thymus Expressed  
ChemoKine. TECK has weak homology with other CC chemokines  
and maps to mouse chromosome 8. Besides the thymus, mRNA  
encoding TECK was detected at substantial levels in the  
small intestine and at low levels in the liver. The source  
25 of TECK in the thymus was determined to be thymic dendritic  
cells, while in contrast bone marrow-derived dendritic  
cells do not express TECK. The murine TECK recombinant  
protein showed chemotactic activity for activated  
macrophages, dendritic cells and thymocytes. We conclude  
30 that TECK represents a novel thymic dendritic cell-specific  
CC chemokine which is possibly involved in T-cell  
development.

Chemokines belong to a family of small peptides (6-15  
kDa) whose best described biological function is to control  
35 the migration of certain leukocyte populations to localized  
sites of inflammation. Baggiolini, et al. (1994) Adv. in  
Immun. 55:97-179; Schall and Bacon (1994) Curr Opin Immun

6:865-873; Hedrick and Zlotnik (1996) Curr. Opin. Immunol.  
8:343-347. In the last few years many new members of the  
chemokine super family have been characterized.  
Initially, new chemokines were identified through their  
5 chemotactic effects on leukocytes (Baggiolini et al.  
(1994); Schall and Bacon (1994)) and were isolated mainly  
from blood leukocytes or cell lines. More recently,  
approaches based on the selective cloning of secreted  
molecules by signal sequence trap (Tashiro, et al. (1993)  
10 Science 261:600-603; Imai, et al. (1996) J. Biol. Chem.  
271:21514-21521) or on the exploitation of public and  
private databases of expressed sequence tags (EST) through  
bioinformatics (Hieshima, et al. (1997) J. Biol. Chem.  
272:5846-5853; Patel, et al. (1997) J. Exp. Med. 185:1163-  
15 1172; and Rossi, et al. (1997) J. Immunol. 158:1033-1036),  
have allowed the rapid identification of novel chemokines  
based on sequence and structural homologies. These  
approaches take advantage of the fact that most of the  
chemokines are secreted factors whose protein sequence  
20 contain four conserved cysteines (Schall (1994) "The  
Chemokines" pp. 419-460 in Thomson (eds.) The Cytokine  
Handbook, Academic Press, New York. The CXC or  $\alpha$  chemokine  
family has the two first amino-terminal cysteines separated  
by a non-conserved amino acid. In the CC or  $\beta$  chemokine  
25 family, these two cysteines are consecutive. A third type  
of chemokine, the C or  $\gamma$  family, is represented by  
lymphotactin, which conserves two cysteines (1 and 3)  
instead of the original four (Kelner, et al. (1994) Science  
266:1395-1399). Finally, a recently identified chemokine  
30 with three amino acids separating the first two cysteines  
defines a fourth CX<sub>3</sub>C family (Bazan, et al. (1997) Nature  
385:640-644).

Interestingly, some of the new chemokines discovered  
show a relatively restricted pattern of expression (Imai et  
35 al. (1996); Hieshima et al. (1997)). It is tempting to  
suggest that these new approaches may lead to the discovery  
of tissue- or cell-specific chemokines. In addition, new



biological evidence for important new roles of chemokines in hemopoiesis (Cook (1996) J. Leukoc. Biol. 59:61-66; and Nagasawa, et al. (1996) Nature 382:635-638) and the control of viral infections including HIV (Cocchi, et al. (1995) Science 270:1811-1815; and Cook, et al. (1995) Science 269:1583-1585). Thus, the molecular cloning of novel chemokines through DNA-based strategies may uncover novel proteins belonging to the chemokine super family but whose physiological role goes beyond the control of inflammation.

10 In an attempt to identify novel genes involved in T-cell development, we analyzed a cDNA library from the thymus of Recombinase Activation Gene-1 (RAG-1) deficient mice. We identified a novel CC chemokine designated TECK for Thymus Expressed ChemoKine, based on sequence homology with other known chemokines. We subsequently isolated the human homologue of TECK. The pattern of expression of TECK mRNA is highly restricted to the thymus and small intestine in both human and mouse. Moreover, in the mouse thymus, TECK protein is produced by dendritic cells while splenic dendritic cells do not express TECK mRNA. Recombinant TECK showed chemotactic activity on thymocytes, macrophages, THP-1 cells and dendritic cells, while it was inactive on peripheral lymphocytes and neutrophils. The restricted pattern of expression of TECK together with its biological properties suggest a role for this novel dendritic cell-specific chemokine in T-cell development.

#### A. Cloning and structural analysis of mouse TECK

A directional cDNA library was made from RAG-1 deficient mouse thymus and analyzed by random sequencing. One of the clones contained an open reading frame with significant homology to previously described CC chemokines. The full-length cDNA contains 1037 bp including an open reading frame of 426 bp encoding a protein of 142 amino acids and will be identified in this report as mTECK (see Table 1). In the 3' untranslated region, there is one unique polyadenylation signal consistent with the single mRNA species observed in northern blots. The mTECK cDNA

does not contain any ATTTA transcript destabilization motif (Shaw and Kamrn (1986) Cell 46:659-667). The comparison of the amino acid sequence of mTECK with previously described murine CC chemokines demonstrates the conservation of the four cysteines present in all these chemokines. However, mTECK shows few additional identities with these proteins.

#### B. Cloning and molecular characterization of human TECK

To investigate the possible existence of a gene homologous to mTECK in other mammalian species, a Southern blot with genomic DNA from various species was hybridized with the mTECK cDNA probe. Under high stringency conditions, hybridizing bands were detected in mouse, rat, hamster and human genomic DNAs. Interestingly, a single band was detected in human, suggesting that a single gene encodes for TECK in this species. The multiple bands present in mouse, rat and hamster could be the result of a internal EcoRI site within the TECK gene. Alternatively, the TECK gene may have been duplicated in these species.

In order to clone the human homologue of mTECK, a blot of cDNAs from a panel of human cDNA libraries was hybridized with the mTECK cDNA probe. A signal was observed in a fetal small intestine cDNA library.

Screening of this library with the mTECK probe allowed the isolation of several identical clones of 1012 bp with an open reading frame of 453 bp encoding a protein of 151 amino acids. This protein had a much higher degree of homology at the nucleic acid level (71% nucleic acid identity for the open reading frame and 49.3% amino acid identity) to mTECK than to other known CC chemokines and was thus designated as hTECK.

#### C. Chromosomal location of mTECK

It has been shown that the genes encoding for most chemokines are clustered in the genome. The genes encoding CC chemokines cluster on mouse chromosome 11 and human chromosome 17q11-12. Schall (1994); and Hedrick and Zlotnik (1996). The chromosomal location of mTECK (designated gene symbol Teck) was determined by inter

specific backcross analysis between ([C57Bl/6J X M.  
spretus]F1 X C57Bl/6J) mice. Jenkins, et al. (1982) J.  
Virol. 43:26-36. The mapping results indicated that the  
Teck locus is located on the proximal part of mouse  
5 chromosome 8. Although the chromosomal location of the  
human Teck locus could not be determined, this region of  
mouse chromosome 8 is syntenic to the human 19p13.3 and  
13q34 regions. However, the Teck locus is also very close  
to a region syntenic to human chromosome 4. The closest  
10 known gene, Insr, encodes the insulin receptor and the  
genetic distance between Teck and Insr was estimated at  $0.9$   
 $\pm 0.9$  cM. We have compared our inter specific map of  
chromosome 8 with a composite mouse linkage map that  
reports the map location of many uncloned mouse mutations  
15 (Mouse Genome Database, The Jackson Laboratory, Bar Harbor,  
ME). Teck resides in a region of the composite map that  
lacks mouse mutations with a phenotype that might be  
expected for an alteration in this locus.

D. Analysis of mTECK and hTECK mRNA distribution in cells  
20 and tissues

An analysis of the distribution of mTECK mRNA in  
tissues and cells by northern blotting or by Southern  
blotting of mouse cDNA libraries revealed that mTECK was  
expressed at significant levels only in the thymus and to a  
25 lesser extent in small intestine (Table 7). Weak  
expression of mTECK mRNA was observed in brain, testis, and  
liver RAG-1-/- cDNA libraries. Interestingly, mTECK mRNA  
was detected in a cDNA library of activated pro-T cells.  
Pro-T cells represent an early stage of intra-thymic T-cell  
30 progenitors, not fully committed to the T-cell lineage  
since they can give rise to NK and dendritic cells. Moore  
and Zlotnik (1995) Blood 86:1850-1860; Wu, et al. (1996).  
J. Exp. Med. 184:903-911). In contrast, mTECK mRNA was  
undetectable in resting or activated thymocytes, peripheral  
35 T or B cells, macrophages, PBLs, splenic dendritic cells  
and in all other tissues tested, with the exception of  
spleens recovered from mice injected with LPS (Table 7).

Interestingly, mTECK mRNA was detected by PCR in fetal thymuses of day 14 of gestation, indicating that mTECK is expressed in the thymus at the earliest stages of T-cell development.

5       The distribution of hTECK mRNA was similarly analyzed. As with mouse, hTECK mRNA expression was highly restricted to the thymus and small intestine. Weak expression was also detected in inflamed tonsil and fetal spleen, but at much lower levels than that observed in the thymus since  
10       this particular blot was exposed for a long time. Importantly, hTECK mRNA was absent from a series of cDNA libraries from dendritic cells derived in vitro from bone marrow CD34+ progenitors cells or peripheral blood monocytes. In addition, hTECK mRNA was also absent from  
15       libraries of monocyte-derived dendritic cells stimulated with LPS or a combination of TNF- $\alpha$ , IL-1 $\alpha$  and monocyte supernatant for 4 and 16 hours. Collectively, these data indicate that TECK mRNA is specifically expressed at high levels in thymus and small intestine in vivo.

20       E. Identification of mTECK-producing cells in vivo

      The abundance of mTECK mRNA expression in RAG-1 deficient thymus and its absence in thymic T cells suggested that mTECK was expressed by a thymic stromal component in normal mice. We performed in situ mRNA  
25       hybridization with sense or antisense mTECK probes generated by PCR. Thymic sections hybridized with the sense probe (negative control) demonstrated no specific staining while sections hybridized with the anti-sense probe at the same concentration showed specific staining in  
30       the thymic medulla. At higher magnification, positive cells appeared to have a non-lymphoid morphology with processes surrounding lymphoid cells. This experiment indicated that, in vivo, mTECK mRNA is expressed by a non-lymphoid component of the medullary stroma, possibly  
35       dendritic cells.

      The thymic stroma is mainly composed of epithelial cells, macrophages, dendritic cells and fibroblasts,

together with a network of vascular and nervous tissue. Boyd, et al. (1993) Immunol. Today 14:445-459. Since we previously failed to detect mTECK mRNA expression in thymic epithelial or macrophage cell lines with or without  
5 activation with IFN- $\gamma$  (Table 7), we sorted thymic dendritic cells based on their high expression of MHC class II and CD11c (N-418 antibody). Analysis of mTECK expression by RT-PCR revealed that freshly isolated MHC class II+ CD11c+ thymic dendritic cells expressed mTECK mRNA while the MHC  
10 class II+ CD11c- subset was negative. In contrast, mTECK mRNA was undetectable in a cDNA library made from freshly isolated splenic dendritic cells (Table 7).

We then performed immunostaining of thymic sections and purified thymic dendritic cells with a polyclonal  
15 antibody raised against a decapeptide corresponding to the C-terminus of mTECK. This polyclonal antibody reacted with recombinant mTECK in ELISA and western blot while normal rabbit serum was negative. In thymic sections, the polyclonal anti-peptide antibody reacted with a stromal  
20 component of the thymic medulla consistent with the in situ hybridization data while staining with normal rabbit serum was negative. Interestingly, the antibody also reacted weakly with some endothelial cells, raising the possibility that mTECK can be produced by the thymic endothelium.

25 Finally, the anti-mTECK polyclonal antibody stained sorted thymic dendritic cells, while the control serum was negative. High magnification clearly showed intra-cellular staining of cells with characteristic dendritic morphology. Taken together, these results indicate that thymic  
30 dendritic cells and possibly thymic endothelial cells are producing TECK in vivo.

#### F. Chemotactic activities of mTECK protein

To evaluate the biologic properties of mTECK, a recombinant protein with a N-terminal FLAG peptide was  
35 obtained in a bacterial expression system. In some experiments, a recombinant mTECK protein with a C-terminal FLAG was used and similar results were obtained.

Interestingly, mTECK induced the migration of mouse thymocytes (Figure 1A). The optimal response was obtained with a dose of 10 ng/ml TECK. Cell migration was determined to be chemotaxis and not chemokinesis through the checkerboard analysis. Furthermore, it is established that chemokines bind to specific receptors that are coupled through heterotrimeric G proteins to intra-cellular signal-transducing pathways. Murphy (1994) Annu. Rev. Immunol. 12:593-633. To determine whether the chemotaxis of thymocytes involved a G protein-coupled receptor, cells were incubated prior to the assay with 10 ng/ml pertussis toxin which ADP-ribosylates G $\alpha_i$ -proteins. Katz, et al. (1992) Nature 360:686-689. This pre-treatment completely abrogated the chemotactic response of thymocytes to mTECK (Figure 1A).

The recombinant mTECK protein also induced the migration of human monocytic THP-1 cells activated for 16 hours with IFN- $\gamma$  (Figure 1B), while it was not significantly active on resting THP-1 cells. This experiment showed that mTECK is active on human cells. In addition, mTECK induced activated mouse peritoneal macrophages to migrate as well as highly purified mouse splenic dendritic cells (Figure 1B). In all these experiments, the optimal dose of mTECK was 10 ng/ml. In contrast, no chemotaxis was observed with bone marrow cells, purified neutrophils, splenic B cells, splenic T cells or IL-2 activated RAG-1 deficient mouse splenocytes lacking mature T and B lymphocytes (Mombaerts, et al. (1992) Cell 68:869-877) and therefore enriched in NK cells. These data are consistent with the absence of in vivo accumulation of neutrophils, monocytes or lymphocytes 2 and 5 h following an intra-peritoneal injection of 10  $\mu$ g mTECK. Collectively, these data indicate that TECK is a chemotactic factor for thymocytes, macrophages and dendritic cells.

G. TECK, a distant member of the CC chemokine family

In this report, we describe the molecular isolation and characterization of TECK, a novel mouse and human CC

chemokine. Analysis of its predicted amino acid sequence showed that TECK is distantly related to previously described CC chemokines. Conservation of particular amino acids among most CC chemokines may be related to their functional importance. Beall, et al. (1992) J. Biol. Chem. 267:3455-3459; and Lusti-Narasimhan, et al. (1995) J. Biol. Chem. 270:2716-2721. In particular, a tyrosine residue between the second and third cysteines has been shown to be critical for monocyte chemotaxis (in position 50) (Beall et al. (1992)). While TECK does not have a tyrosine at this particular position, it has one in position 52 that may have the same function, since TECK is chemotactic for activated monocytes. In addition to these differences in the primary structure, the gene encoding TECK maps on chromosome 8 in the mouse, unlike most other CC chemokines which are clustered on chromosome 11. This is not the first report of an unusual chromosomal location for a CC chemokine. We have cloned the human CC chemokine MIP-3 $\beta$  and showed that its gene was on chromosome 9 rather than 17 (Rossi, et al. (1997)), and the gene encoding the novel human CC chemokine MIP-3 $\alpha$ /LARC (Rossi, et al. (1997)) has been mapped on chromosome 2 (Hieshima, et al. (1997)). It is likely that the CC chemokines on chromosome 11 in the mouse and 17 in human have been generated through gene duplication of a primordial chemokine. Our results suggest that TECK may have been generated at an earlier stage during evolution. In this regard, the TECK gene may have evolved to ensure functions similar to other CC chemokines with a distant primary structure but through similar receptor(s) as dictated by its secondary and tertiary structures. Alternatively, the receptor(s) and physiological role of TECK may be unique among chemokines. H. TECK expression and function is associated with T-cell development

We observed that TECK was strongly expressed in the thymus which is the primary lymphoid organ where T-cell development takes place. Recently, another CC chemokine

highly expressed in the thymus, TARC, has been identified. Imai, et al. (1996). However, TARC is also expressed in lung and colon as well as activated PBMC (Imai, et al. (1996)) while TECK was absent from these tissues. Besides  
5 the thymus, numerous reports indicate that T cell development can occur in the small intestine (Poussier and Julius (1994) Annu. Rev. Immunol. 12:521-553) where TECK is also expressed. Interestingly, the liver has also been suggested to support T-cell development to some extent  
10 (Abo, et al. (1994) Int. Rev. Immunol. 11:61-102) and we observed a low TECK expression in a liver cDNA library. These data show that TECK expression correlates with organs that support T-cell development.

While many molecular and cellular aspects of T-cell  
15 differentiation are well documented, the precise role of chemokines in T-cell development is still unknown. Recently, it has been shown that the bone marrow stroma-derived CXC chemokine SDF-1 is important for B lymphopoiesis and myelopoiesis since SDF-1  $-/-$  mice are  
20 impaired for these functions (Nagasawa, et al. (1996)). Similarly, it is likely that chemokines act at different steps of T-cell differentiation. Chemokines, together with the expression of appropriate adhesion molecules, may dictate the migration of uncommitted progenitors from the  
25 bone marrow to other anatomic locations. Indeed SDF-1 is chemoattractant for human CD34+ progenitor cells. Aiuti, et al. (1997) J. Exp. Med. 185:111-120. The observation that TECK is chemoattractant for thymocytes but not for mature peripheral T cells suggests that TECK could attract  
30 T-cell progenitors to the thymus. Such populations are very difficult to isolate in sufficient numbers to conduct in vitro chemotaxis experiments, but we are currently designing new strategies to address this important question. In addition, we have not found significant  
35 chemotactic activity of TECK on bone marrow cells. SDF-1 was shown to be much less potent on CD34+ progenitors from the peripheral blood than those from the bone marrow.



Aiuti, et al. (1997). It is possible that the sensitivity of progenitor cells to TECK would increase as these cells leave the bone marrow to colonize lymphoid organs.

Importantly, intra-thymic maturation is also characterized by a directional migration from the subcapsular region which contains the earliest progenitors to the cortex and finally to the medulla where thymocytes finish their maturation (Boyd, et al. (1993)). It is possible that the secretion of TECK by medullary dendritic cells may play a role in this directional migration. Yet another possibility is that TECK may play a role in the organization and development of the thymic stroma.

We also showed that TECK is chemotactic for activated macrophages and dendritic cells. These two cell types also play important roles in T-cell development. Through a complex screening process involving positive and negative selection events most of the antigenic specificities randomly generated in the thymus will be eliminated by programmed cell death (Janeway (1994) Immunity 1:3-6). The efficient scavenging of dead thymocytes is probably mediated, at least partially, by thymic macrophages and thus TECK could play an important role through its action on activated macrophages. Further along, T-cells with a high affinity for self-antigens and thus potentially harmful are eliminated through negative selection (Janeway (1994)). It is believed that thymic dendritic cells are primarily responsible for the negative selection of thymocytes, therefore playing a major role in the establishment of tolerance. Inaba, et al. (1991) J. Exp. Med. 173:549-559. An efficient mechanism of central tolerance should eliminate T cells potentially reactive against auto-antigens which are not expressed in the thymus, such as organ specific auto-antigens. Several known chemokines induce the migration of dendritic cells and could therefore contribute to their recruitment during peripheral immune responses. Sozzani, et al. (1995) J. Immunol. 155:3292-3295; and Xu, et al. (1996) J. Leukoc.

Biol. 60:365-371. Similarly, dendritic cells presenting organ-specific or other antigens could be recruited to the thymus or the small intestine and induce negative selection of T cells specific for these antigens. It is possible that thymus- and small intestine-specific chemokines active on dendritic cells such as TECK could play an important role in the establishment of tolerance. Thus, TECK could potentially interact at several important steps of T-cell development. Future experiments will aim to define the precise role of TECK in T-cell development and other physiological processes through the use of genetically modified mice.

I. TECK is specifically expressed by thymic dendritic cells

Dendritic cells represent an heterogeneous cell population derived from bone marrow progenitors. They are present in non-lymphoid organs as immature dendritic cells (such as Langerhans cells in the skin) where they display a high ability for antigen capture. Cella, et al. (1997) Curr. Opin. Immunol. 9:10-16. Subsequent to antigen challenge, they will migrate to secondary lymphoid organs and will acquire a high capacity to present processed antigens to naive T-cells to initiate a specific immune response (Cella, et al. (1997)). It has been shown that dendritic cells can derive from CD34+ progenitors cultured in the presence of GM-CSF and TNF- $\alpha$  (Caux, et al. (1992) Nature 360:258-261; and Caux, et al. (1996) J. Exp. Med. 184:695-706) or from monocytes in the presence of GM-CSF and IL-4 (Sallusto and Lanzavecchia (1994) J. Exp. Med. 179:1109-1118). Interestingly, there is also evidence for a lymphoid dendritic cell precursor in thymus and bone marrow which is able to derive both lymphocytes and dendritic cells in the absence of GM-CSF. Ardavin, et al. (1993) Nature 362:761-763; Galy, et al. (1995) Immunity 3:459-473; Marquez, et al. (1995) J. Exp. Med. 181:475-483; and Wu, et al. (1996). These lymphoid-derived dendritic cells may have different functional properties such as a

negative regulation of T-cell responses since they express FasL in the mouse. Suss and Shortman (1996) J. Exp. Med. 183:1789-1796. We found that TECK was expressed at high levels in mouse thymic dendritic cells but was absent in cDNA libraries from mouse splenic dendritic cells or from human dendritic cells generated in vitro from CD34+ precursors or monocytes. Interestingly, mTECK mRNA was present at a low level in a population of early thymocyte progenitors still able to derive dendritic cells (Wu, et al. (1996). Thus, it would be tempting to suggest that TECK could be a specific marker of lymphoid-derived dendritic cells. However, we observed that TECK was absent from splenic dendritic cells that likely contain lymphoid-derived dendritic cells. The expression of TECK mRNA appeared in the spleen of mice injected with LPS would suggest that peripheral dendritic cells may express TECK upon activation, but we found that TECK was not expressed in cDNA libraries of bone-marrow derived dendritic cells activated with LPS, PMA and ionomycin or IL-1 $\alpha$  and TNF- $\alpha$ .

It is possible that the normal expression of TECK is specific for lymphoid-derived dendritic cells or, alternatively, that it is upregulated by very specific stimuli present in the thymic and intestinal micro-environment under physiological conditions. Consistent with the latter hypothesis is our observation of specific staining of thymic endothelial cells with anti-TECK antibody since we have not been able to find TECK expression in human HUVEC endothelial cells by northern blot analysis, without activation or following a 16 hour-activation with various combinations of IL-1, TNF- $\alpha$ , IL-4, IL-7 and oncostatin while some of these stimuli induce the expression of other CC chemokines in endothelial cells. Rollins and Pober (1991) Am. J. Pathol. 138:1315-1319; Marfaing-Koka, et al. (1995) J. Immunol. 154:1870-1878; Garcia-Zepeda, et al. (1996) J. Immunol. 157:5613-5626; and Garcia-Zepeda, et al. (1996) Nat. Med. 4:449-456. Taken together, our data strongly suggest that TECK is a novel

chemokine specifically expressed by activated lymphoid-derived dendritic cells.

Through their function of antigen presentation, dendritic cells play major roles in the establishment of tolerance and in the initiation of an antigen-specific immune response. The use of purified dendritic cells has been recently proposed in different therapeutic protocols (Cella, et al. (1997)). The discovery of factors with a regulated expression in dendritic cells such as the novel CC chemokine TECK will certainly improve our knowledge of the biology of dendritic cells and lead to the design of relevant in vivo applications.

#### J. Mice and in vivo experimental procedures

Four to eight week-old and time-pregnant BALB/c mice were purchased from Simonsen Laboratories (Gilroy, CA). RAG-1-deficient mice (Mombaerts, et al. (1992)) were purchased from The Jackson Laboratory (Bar Harbor, ME). To analyze TECK expression after in vivo activation, various organs were recovered from pools of 2 mice 3 hours after intravenous LPS injection (50 µg LPS in 200 µl PBS or 200 µl PBS for controls).

#### K. Cell purification, culture and stimulation.

THP-1 cells (TIB-202 from the American Type Culture Collection, Rockville, MD) were cultured in complete medium which consisted in RPMI 1640 medium (JRH BioSciences, Lenexa, KS) supplemented with 10% FCS, 200 mM L-glutamin,  $5 \times 10^{-5}$  M mercaptoethanol, MEM amino-acids and vitamins, sodium bicarbonate, penicillin, streptomycin (all from Sigma, ST. Louis, MO), and gentamycin (Boehringer, Indianapolis, IN). To obtain activated mouse macrophages, 10 ml of cold PBS were injected into the peritoneum and the collected cells allowed to adhere to plastic for 24 h in complete medium. The adherent fraction, mostly macrophages, was then collected. To obtain splenic dendritic cells, a splenocyte cell suspension was prepared in RPMI 1640 Dutch modified medium (Life Technologies,

Paisley, Scotland) as described previously in, e.g.,  
Macatonia, et al. (1987) J. Exp. Med. 166:1654-1667.

Splenocytes were incubated at 37° C for 16 h and the cell  
suspension was collected and laid over Metrizamide (Nycomed  
5 Pharma, Oslo, Norway). After centrifugation for 10 min. at  
1700 x g, the low interface was collected and stained with  
anti-Mac-1 (Pharmingen, San Diego, CA) and the anti-CD11c  
N-418 antibodies (Macatonia, et al. (1993) J. Immunol.  
150:3755-3765). Splenic dendritic cells were sorted by  
10 flow cytometry on a FACStar plus cell sorter (Becton  
Dickinson, Mountain View, CA) to a purity greater than 98%  
upon reanalysis in all the experiments included in this  
report. To obtain thymic dendritic cells, thymuses were  
cut in small fragments and resuspended in 10 ml of RPMI-  
15 1640 +10% FCS containing 1 mg/ml collagenase and 0.02 mg/ml  
DNase I (both from SIGMA) and digested with continuous  
agitation at room temperature for 30 min. (Shortman, et  
al. (1995) Adv. Exp. Med. Biol. 378:21-29). One ml of 0.1M  
EDTA pH 7.2 was added for an additional 5 min. Cells were  
20 then washed in complete medium, resuspended in complete  
medium and overlaid onto Metrizamide. The thymic dendritic  
cell-enriched preparation was then stained with anti-IAD  
and N-418 antibodies and the dendritic cells sorted by flow  
cytometry

## 25 L. Molecular cloning of mouse and human TECK

The cDNA encoding mouse TECK was obtained by random  
sequencing of a RAG-1 KO mouse thymic directional cDNA  
library. Briefly, mRNA was extracted using RNazol™ B  
(Tel-Test, Friendswood, TX) and then oligotex-dT mRNA kit  
30 (Quiagen, Chatsworth, CA) following the manufacturer's  
instruction. A directional cDNA library was prepared using  
the Superscript™ Plasmid System (Gibco-BRL, Grand Island,  
NY) and cloned into the pME18s plasmid vector. Sequencing  
was done using the TaQ DyeDeoxy Terminator Cycle Sequencing  
35 kit (Applied Biosystems, Foster City, CA). To determine  
whether TECK was present in other mammals including human,  
a Southern blot containing EcoRI digested genomic DNA from

different species (Bios Laboratories, New Haven, CT) was hybridized with the full-length mouse TECK cDNA.

The cDNA encoding human TECK was found by screening of a small intestine cDNA library using the full-length mouse TECK cDNA as a probe following standard procedures.

M. Northern blot analysis of RNA and Southern blot analysis of cDNA libraries

All RNA's were isolated from tissues or cells using RNazol™ B (Tel-Test) and analyzed after electrophoresis in a 1% formaldehyde-agarose gel (10 µg/lane). RNA's were then blotted onto a Hybond-N+ nylon membrane (Amersham, Arlington Heights, IL). Some northern blots of mRNA were bought from Clontech (Palo Alto, CA). To analyze the expression of TECK in cDNA libraries (obtained from T. MacClanahan, DNAX), 10 µg of DNA were digested with the appropriate restriction enzymes to release their inserts and analyze by Southern blotting onto nylon membranes. Northern blots and blots of cDNA libraries were hybridized for 16 hours at 65°C with a <sup>32</sup>P-labeled probe consisting in the full-length cDNA encoding for mouse or human TECK and then washed and exposed, according to standard protocols.

N. Inter specific mouse backcross mapping

Inter specific backcross progeny were generated by mating (C57Bl/6J x M. spretus) F1 females and C57Bl/6J males as described, e.g., in Copeland and Jenkins (1991) Trends Genet. 7:113-118. A total of 205 N<sub>2</sub> mice were used to map the Teck locus. DNA isolation, restriction enzyme digestion, agarose gel electrophoresis, Southern blot transfer and hybridization with the full-length mTECK cDNA probe were performed as described, e.g., in Jenkins, et al. (1982). Fragments of 7.5, 6.9, and 2.5 kb were detected in HincII digested C57Bl/6J DNA and fragments of 8.8 and 5.4 kb were detected in HincII digested M. spretus DNA. The presence or absence of the 8.8 and 5.4 kb HincII M. spretus-specific fragments, which cosegregated, was followed in backcross mice. A description of the probes and RFLPs for two of the loci linked to Teck including Insr

has been reported previously, e.g., in Ceci, et al. (1990) Genomics 6:72-79. Recombination distances were calculated as described (Green (1981) "Linkage, recombination and mapping" pp. 77-113 in Genetics and Probability in Animal Breeding Experiments, Oxford University Press, New York) using the computer program SPRETUS MADNESS.

#### O. Measurement of TECK mRNA expression by RT-PCR

RNA's from sorted thymic dendritic cells or fetal thymuses were prepared with the RNeasy total RNA kit (Quiagen, Chatsworth, CA), following the manufacturer's instructions. First strand cDNAs were generated by reverse transcription with a random hexamer in a 10 µl reaction and 1 µl of this reaction was used as a template for PCR. TECK expression was compared to the expression of hypoxanthine-guanine phosphoribosyl transferase (HPRT). Primer sequences were as follows: TECK: 5' primer, 5'CCTTCAGGTATCTGGAGAGGAGATC3' (nucleotides 58-72 of SEQ ID NO: 1) and 3' primer, 5'CACGCTTGTTGTTGGGGTTC3' (complement of nucleotides 447-468 of SEQ ID NO: 1), HPRT: 5' primer, 5'GTAATGATCAGTCAACGGGGGAC3' (SEQ ID NO: 17) and 3' primer, 5'CCAGCAAGCTTGCAACCTTAACCA3' (SEQ ID NO: 18). Samples were submitted to 25 cycles of amplification, each composed of 94° C for 1 min., 57° C for 30 s and 72° C for 2 min. PCR products were then separated by electrophoresis in 2% agarose gels and stained with ethidium bromide.

#### P. In Situ Hybridization

Biotin-14-CTP labeled sense and antisense riboprobes were generated using a non radioactive RNA labeling system (Gibco, Gaithersburg, MD) and the plasmid PCR11 (InVitrogen, Carlsbad, CA) containing a 400 base pair TECK cDNA fragment inserted by PCR and TA cloning (InVitrogen). Paraffin-embedded tissues were cut in 3-5 µm sections, mounted on slides, baked at 60° C for one hour, deparaffinized in xylene (Fisher Scientific, Pittsburgh, PA) and immersed in 100% ethanol. Sections were then incubated for 10 min at 37° C in proteinase K solution (40 mg/ml) (Gibco) in PBS and rinsed for 2 min in PBS at room

temperature before being refixed in 10% formalin (Fisher Scientific, Pittsburgh, PA) in PBS for 1 min. Next, the sections were dehydrated through graded solutions of ethanol and air dried. Hybridization was carried out using the Gibco in situ hybridization and detection system kit. Vanadyl ribonucleoside complex (Gibco) was added to the hybridization solution (39 mM final). A 0.1 µg/ml concentration of each probe was used during an 18 h hybridization at 42° C. Post-hybridization washes used room temperature 0.2X SSC. Following detection and substrate visualization, the slides were counterstained with 1% nuclear red stain (Sigma, St. Louis, MO).

#### Q. Immunohistochemistry

A polyclonal antibody specific of a synthetic decapeptide identical to the C-terminus part of murine TECK (Figure 1) was prepared in rabbits by Research Genetics (Huntsville, AL). Normal rabbit serum from a pool of 50 different animals (Research Genetics) was used as a negative control. To study TECK protein expression in the mouse thymus, 6 µm thick cryostat sections were thaw mounted on organosilicone subbed slides (American Histology Reagent Co., Stockton, CA.) and fixed in 3% formalin (Fisher Scientific, Springfield, NJ) in Hank's Balanced Salt Solution with 0.01M HEPES (HBSS-HEPES), pH 7.4, for 15 min at room temperature. The sections were sequentially blocked for endogenous biotin binding using the Vector blocking kit (Vector Laboratories, Burlingame, CA) and for endogenous peroxidase activity with a 1% hydrogen peroxide, 0.2M sodium azide solution, in HBSS-HEPES with 0.1% saponin (staining buffer). Non-specific antibody binding sites were then blocked with 10% normal goat serum (Sigma) in staining buffer. Sections prepared as above were first incubated for 18 h at 25°C with 1/500 dilution of polyclonal antibody or control rabbit serum in staining buffer. In the second step, the sections were incubated for 1 h at room temperature with biotin labeled goat anti-rabbit IgG (2 µg/ml) (Vector Laboratories) in staining



buffer and then for 30 min at room temperature with the Vectastain Elite ABC Kit (Vector Laboratories) in staining buffer. The sections were then rinsed in HBSS-HEPES without saponin. Immunoenzyme tissue staining was revealed with 3, 3' - diaminobenzidine tetrahydrochloride (DAB) substrate (0.5 mg/ml) (Sigma) in 0.05M Tris, pH 7.4, containing 0.0075% hydrogen peroxide. The substrate reaction was stopped by rinsing the sections in distilled water. The sections were then counterstained with Harris' hematoxylin (Shandon Lipshaw, Pittsburg, PA).

The expression of TECK mRNA in murine adult thymus was analyzed by in situ hybridization and revealed a discrete positive non-lymphoid population within the thymus medulla. The expression of TECK protein was analyzed by using a polyclonal anti-serum made in a rabbit immunised with a peptide that consisted in the last 12 amino-acid of the murine TECK protein sequence. This polyclonal antibody reacts with the murine TECK recombinant protein prepared at DNAX both in ELISA and western blot. The application of this anti-serum on mouse adult thymic sections confirmed the distribution pattern obtained by in situ hybridization: the cells producing TECK are medullary stromal cells. The precise cell type producing TECK within the mouse thymus was identified, using the same anti-serum on sorted thymic subsets, as being the thymic dendritic cells.

R. Production of recombinant mouse TECK in *Escherichia coli* and other chemokines

Mouse recombinant TECK was produced in *E. coli* as a N-terminal FLAG (DYKDDDDKL; SEQ ID NO: 19) fusion protein. Briefly, the fusion construct containing FLAG followed by the mTECK sequence minus the leader peptide (see Table 1) was obtained by PCR amplification of the TECK cDNA in order to flank the coding sequence with HindIII and EcoRI sites and subsequent ligation in the pFLAG.1 vector which contains the FLAG sequence and an OmpA signal sequence. Electro-competent UT 4400 *E. coli* were transformed with the pFLAG.1-mTECK plasmid. The cells were grown in 2 x LB plus

50 µg/ml Ampicillin, induced at an OD. of 2.3 with 400 µM IPTG and harvested. The cell pellet was resuspended in cold lysis buffer (20 mM Tris pH 8, 2 mM EDTA, 20% sucrose, 0.1 mg/ml lysozyme, 100 µl Benzonase), homogenized and allowed to sit for 30 min. Then the same amount of a 1:4 dilution of cold lysis buffer without lysozyme was added for 10 more min. The solution was spun and the supernatant was filtered through a 0.2 µm membrane and then diluted 1:1 in 50 mM Tris pH 7.5. The diluted osmotic extract was submitted to chromatography on a Q-sepharose column equilibrated with 50 mM Tris pH 7.5 and eluted with a linear salt gradient. The fractions containing the recombinant protein were pooled. The fractions were then loaded onto a S-sepharose column equilibrated with 20 mM acetate pH 4.0. The column was eluted with a linear salt gradient and then with a 1.5M NaCl wash that contained the protein. Finally, the eluate was loaded onto a reverse phase column. The column was eluted with a linear gradient of 20% to 80% acetonitrile + 0.1% TFA. The concentration of the mTECK protein was estimated by Comassie blue staining and densitometric scanning of a 10% Nu-PAGE gel with lysozyme as a standard. The purity was estimated at 100% by sequencing of the N-terminus of the recombinant protein. Recombinant murine MIP-1α (R&D Systems, Minneapolis, MN) and lymphotactin (Hedrick, et al. (1997) J. Immunol. 158:1533-1540) were used as controls.

#### S. Assay for chemotaxis

The in vitro migration of cells isolated as described above in response to TECK or other factors was assessed in a modified Boyden micro chamber (Neuroprobe, Cabin John, MD) as described previously (Kelner, et al. (1994)). Briefly, factor dilutions in DMEM medium (Gibco) were loaded in the lower compartment in duplicate and 10<sup>5</sup> cells in a 50 µl volume of DMEM were loaded in the upper compartment. The two compartments were separated by a 5-µm or 8-µm pore size polycarbonate filter (Nucleopore, Pleasanton, CA). After incubation at 37° C for 80 min (or

120 min for lymphocytes), the filters were fixed in methanol and stained with Fields A and B. Cell migrated on the other side of the membrane were counted per five high-power fields (100 x) under microscope. The chemotactic index was calculated from the number of cells counted with the test sample divided by the number of cells counted with medium alone.

Northern blot analysis was performed of RNA from different organs hybridized with the mTECK cDNA probe with or without in vivo LPS stimulation. Hybridizing bands corresponded to the predicted  $\approx 1040$  bp size for mTECK mRNA. Significant induction occurred in spleen (with virtually no background), and in thymus and small intestine (both with higher background); no signal was detected in either condition for heart, lung, kidney, or liver.

mTECK mRNA expression was analysed in the mouse fetal thymus. RNA's from fetal thymic lobes were extracted at day 14, 15, 16 and 17 of gestation. Positive RT-PCR signals were detected in each of day 14, 125, 16, and 17 samples.

mTECK mRNA expression in thymic dendritic cells was evaluated. A population enriched in thymic dendritic cells was prepared from 15 pooled adult thymuses.  $>99\%$  pure dendritic cells were then sorted by flow cytometry based on their MHC Class II+ N-418+ phenotype. mTECK mRNA was then analyzed by RT-PCR and a MHC class II+ N-418- population sorted in the same experiment was used as a negative control. The N418+ sample gave a positive signal, while the N418- sample did not.

Expression analysis was performed with hTECK mRNA in different Human Tissues and Cell Types. Southern blots of human cDNA libraries digested with the appropriate restriction enzymes were hybridized with the hTECK cDNA probe. A major band hybridizing corresponding to the predicted length of hTECK mRNA ( $\approx 1040$  bp) was observed with sometimes some other bands that may represent incomplete cDNAs. Positive signals were detected in tonsil, fetal

spleen, and fetal small intestine. No signal was detected in activated (with PMA and ionomycin for 12 h) NK cells, activated (anti-CD40 antibody and IL-4 for 6 and 12 h) splenocytes,  $\gamma\delta$  T cells, activated (with anti-CD3 and PMA for 6, 12, and 24 h) PBMC, fetal testis, C+ (elutriated monocytes cultured with IFN- $\gamma$  and IL-10) monocytes, C- monocytes, 70% pure dendritic cells (CD1 $\alpha$ + dendritic cell population obtained by expansion of CD34+ bone marrow cells with GM-CSF and TNF- $\alpha$  and resting), and DC3 (similar dendritic cell population stimulated with PMA and ionomycin for 1 and 6 h), DC5 (dendritic cells obtained by culturing peripheral blood monocytes in the presence of IL-4 and GM-CSF), U937 (premonocytic cell line), and CD1 $\alpha$  cell lines. Ras KO mouse cDNA again confirmed that the mouse and human genes crosshybridize.

Four independent lines of transgenic mice expressing TECK in the brain have been made. All animals had neurologic disorders. In addition, several of them suffered severe infections. The consequences of TECK could be a direct one on brain cells which nature remains to be identify. Alternatively, since TECK has been shown in vitro to have effects on macrophages and dendritic cells which are critical effectors of immune responses, the overproduction of TECK could lead to distant effects on these cells at sites of infection. These results suggest that the blockade of TECK production in vivo may help to resolve particular pathological processes, in particular infections. The localization suggests a physiological role in immunological responses involving the thymus, or in colon/small intestine or gastrointestinal inflammation, e.g., Crohn's disease or inflammatory bowel disease.

## VII. Specific Characterization of the M/DC CR (CRAM)

Abbreviations: BAC, bacterial artificial chromosome; bp, base pair; CKR, chemokine receptor; EST, expressed sequence tag; GPR, G-protein-linked receptor; PBMC, peripheral blood mononuclear cells; STS, sequence tagged site.

We describe a novel human gene with high homology to CC- or  $\beta$ -chemokine receptors (CKRs). This putative CKR, CRAM, is most similar to human CCR1, with 46% amino acid identity and 65% similarity. CRAM is encoded by at least two alternatively spliced 1.5 and 1.8 kb mRNAs which specify at least two proteins differing by 12 amino acids at the N-terminus (CRAM-A and CRAM-B). CRAM mRNA was detected mainly in lymphoid tissues and expressed in activated monocytes, but not in B- or T-lymphocytes. CRAM mRNA expression was increased upon stimulation with IFN $\gamma$  and LPS but was not detectably inhibited by interleukin-10. CRAM was localized to the  $\beta$ -CKR cluster at chromosome 3p21 and physically linked to the CCR2 and CCR5 genes. In view of its similarity and genomic linkage to  $\beta$ -CKRs and restricted expression pattern, CRAM may play an important role in immune function. The existence of CRAM with alternative N-termini suggests a mechanism for altering ligand specificity and possibly signalling capacity of a single CKR.

Chemokines play critical roles in the chemoattraction and activation of leukocytes (Premack and Schall (1996) Nat Med 2:1174; Murphy (1996) Cytokine Growth Factor Rev 7:47; and Furie and Randolph (1995) Am J Pathol 146:1287), and have been divided into four families, based on the spacing of the first two of (usually) four conserved cysteine residues. The  $\alpha$  chemokines, with a C-X-C motif, include IL-8, MIP-2 $\alpha$ , GRO $\beta$ , and ENA-78. The  $\beta$  chemokines (C-C motif), include MIP-1 $\alpha$ , MCP-1, TARC, and RANTES. Recently, two new chemokine families have been defined by lymphotactin ( $\gamma$ ) and CX<sub>3</sub>Ckine ( $\delta$ ). Lymphotactin has only a

single cysteine residue at the corresponding location for the C-C or C-X-C motif. Kelner and Zlotnik. (1995) J Leukoc Biol 57:778; Kennedy, et al. (1995) J Immunol 155:203. CX<sub>3</sub>Ckine contains two cysteines separated by three  
5 intervening amino acids, and is tethered to the cell membrane via a long carboxy-terminal tail of mucin-like repeats. Bazan, et al. (1997) Nature 385:640.

Receptors for chemokines (CKRs) are G-protein coupled receptors (GPRs) with seven transmembrane domains. Novel  
10 CKRs have been identified by expression cloning of receptors binding a particular chemokine ligand (Holmes, et al. (1991) Science 253:1280) or mediating HIV fusion (Feng, et al. (1996) Science 272:872), by PCR using degenerate  
primers specific for conserved regions (Meyer, et al. (1996) J Biol Chem 271:14445; Ponath, et al. (1996) J Exp Med 183:2437; Daugherty, et al. (1996) J Exp Med 183:2349; Kurihara and Bravo (1996) J Biol Chem 271:11603; Power, et al. (1995) J Biol Chem 270:19495; Napolitano, et al. (1996) J Immunol 157:2759; and Raport, et al. (1996) J Leukoc Biol  
20 59:18), and by random sequencing efforts followed by sequence analysis. While nearly 30 CKR-like genes have been cloned from mammals and mammalian viruses, only 17 have been shown to bind identified chemokines. Thus, a substantial number of CKR-like molecules remain "orphan  
25 receptors." Most CKRs with experimentally identified ligands bind to more than one ligand. IL-8 receptor B (CXCR2) binds to the  $\alpha$  chemokines IL-8, NAP-2, and MGSA (Suzuki, et al. (1994) J Biol Chem 269:18263), whereas human CCR5 binds the  $\beta$  chemokines RANTES, MIP-1 $\alpha$ , and MIP-  
30 1 $\beta$  (Raport, et al. (1996) J Biol Chem 271:17161; and Alkhatib, et al. (1996) Science 272:1955).

We have used cDNA library subtraction to isolate genes which are induced by monocyte activation. We thereby  
isolated a cDNA clone from a subtracted library enriched  
35 for monocyte activation-specific cDNAs that shows considerable homology to CC- or  $\beta$ -CKRs and maps within the  $\beta$ -CKR cluster on human chromosome 3p21. Expression of this

gene was detected in several lymphoid tissues and in activated monocytes (but not lymphocytes). We provisionally designate this gene CRAM, for chemokine receptor of activated monocytes. CRAM is expressed as at least two alternatively spliced mRNAs encoding CKRs with different N-terminal amino acid sequences, suggesting a possible novel mechanism for regulation of CKR ligand specificity.

#### A. Cell cultures and cDNA library construction

Human PBMC were purified by density gradient centrifugation on Ficoll (Pharmacia Biotech Inc., Piscataway, NJ) using standard procedures. Monocytes were enriched from PBMC by adherence to tissue culture flasks and cultured in DMEM + 10% FCS. Monocytes were activated by culture with 100 ng/ml IFN $\gamma$  (R & D Systems Inc., Minneapolis, MN) and 1  $\mu$ g/ml LPS (Life Technologies, Grand Island, NY) for 1-15 hr. Total RNA was prepared by guanidinium isothiocyanate lysis followed by poly(A)<sup>+</sup> RNA selection using the OLIGOTEX kit (QIAGEN Inc., Chatsworth, CA). cDNA libraries containing  $>2 \times 10^6$  independent clones were constructed using the SuperScript cDNA Kit (Life Technologies).

#### B. cDNA library subtractions

Subtracted cDNA libraries (activated monocytes minus resting PBMC) were constructed. See, e.g., Hara, et al. (1994) Blood 84:189; and Kennedy, et al. (1996) J Interferon Cytokine Res 16:611. The major cDNA species present in the subtracted library were then added (1  $\mu$ g each) to the resting PBMC cDNA library (150  $\mu$ g); this mixture was used as the driver cDNA for a second round of subtraction using 5  $\mu$ g of the activated monocyte cDNA library to enrich for induction-specific cDNAs which were less abundantly expressed.

#### C. DNA sequencing and bioinformatics

The nucleotide sequence of CRAM was determined using an ABI 377 automated sequencer and standard techniques. DNA sequence analyses were performed using Sequencher 3.0

(Gene Codes Corporation, Ann Arbor, MI) and MacVector 6.0 (Oxford Molecular Group). Comparisons to GenBank databases were performed using the BLAST program on web-based servers: (<http://www.ncbi.nlm.nih.gov/BLAST/> and <http://www.genome.ad.jp/SIT/BLAST.html>). Sequence alignments and phylogenetic analyses utilized ClustalW 1.6 (Higgins, et al. (1996) Methods in Enzymology 266:383) and TreeViewPPC 1.2 (Page (1996) Computer Applications in the Biosciences 12:357).

10 D. Analysis of CRAM mRNA expression

Multiple-tissue Northern blots were purchased from Clontech (Palo Alto, CA). Poly(A)+ RNA from human monocytes was used for RNA blot analysis. cDNA libraries from human cells (5 µg) in the pSPORT vector (Life Technologies) were digested with SalI and NotI to release cDNA inserts, electrophoresed on 1% agarose gels, and subjected to Southern blot transfer/hybridization. Hybridizations with <sup>32</sup>P-labeled CRAM DNA fragments encoding the C-terminal 144 amino acids of the predicted ORF were done at 65° C in ExpressHyb (Clontech, Palo Alto, CA) for 2 hr, followed by two stringent washes at 50° C in 0.1X SSC, 0.1% SDS for 45 min. Hybridization was detected using a STORM 860 phosphorimager (Molecular Dynamics, Sunnyvale, CA). Reverse transcriptase PCR (RT-PCR) was performed with Superscript II reverse transcriptase (Life Technologies) and Taq DNA polymerase (Boehringer-Mannheim, Indianapolis, IN). PCR was for 35 cycles of 95° C/45 sec, 62° C/30 sec, 72° C/60 sec. Primers specific for exon 1 (5'-AGACGCTTCAGAGATCCTCTGGAGGCC) or exon 2 (5'-GAAGCTGCTTCGGGGGGTGAGCAAAC) were used in conjunction with an exon 3-specific primer (5'-CAAACACAGCAGAGCAGAGTGATGGCACC) for amplification.

E. Chromosomal localization

PCR was performed on genomic DNA from the 83 cell lines of the Stanford Human Genome Center G3 radiation hybrid panel (Research Genetics, Huntsville, AL) using CRAM primers: (5'-GTGTCCTGGCATGGGTAACAGCC) and



(5'-CGGTGGAATGGTCAGGTTCTTCCC) as previously described for the GeneBridge 4 radiation hybrid panel (Samson, et al. (1996) Genomics 36:522). Data correlating the presence or absence of PCR product to each cell line were entered into the RHserver (Stanford Human Genome Center; <http://shgc.stanford.edu/RH/>). Co-localized STSs were identified on the human physical map using the Entrez server (National Center for Biotechnology Information; <http://www3.ncbi.nlm.nih.gov/Entrez/>).

#### 10 F. cDNA cloning of CRAM

We employed subtractive hybridization to identify genes induced in monocytes upon activation by IFN $\gamma$  and LPS. An activated monocyte cDNA library was first subtracted against a resting PBMC cDNA library. Seven prominent induced cDNAs thus identified were mixed with the resting PBMC library, which was then used as "driver" in another subtraction to generate a new library containing less abundantly expressed, induction-specific cDNAs. More than 100 clones were isolated from this second-round subtracted library, representing 55 unique cDNAs, 25 of which did not correspond to known cDNAs from the non-redundant section of GenBank. One of these clones contained a 1.5 kb insert encoding a large open reading frame with strong homology to all five known human  $\beta$ -CKRs. We designated this cDNA CRAM (chemokine receptor of activated monocytes; or M/DC CR).

#### 25 G. Sequence analysis of CKRs

A phylogenetic analysis of CKRs and related gene sequences revealed two major clades or phylogenetic groups, with several receptors remaining unclustered outside these two groups. Interestingly, the two groups correlated with known ligand specificities: the  $\alpha$ -CKR IL-8RA, IL-8RB, and fusin cluster in a single clade, while  $\beta$ -CKR CCR1 through CCR5 all cluster in a second clade. Of the seven receptors that do not fall into either group, one (DARC) is a promiscuous CKR that binds several  $\alpha$ - and  $\beta$ -chemokines (Neote, et al. (1993) J Biol Chem 268:12247).

The 1536 bp CRAM cDNA encodes an ORF with a predicted size of about 356 amino acids. Phylogenetic analysis showed that CRAM was most closely related to  $\beta$ -CKRs, exhibiting strongest homology to CCR1 (46% identity and 65% similarity), and the least to CCR4, with only 33% identity and 48% similarity. Three other human orphan receptors V28, TER1, and GPR5 also group with  $\beta$ -CKRs, and like CRAM, may be receptors for known or yet to be identified  $\beta$  chemokines.

10 The two most highly conserved regions among CCR1 through CCR5 are in transmembrane region 2 (YLLNLAISDLLF; "TM2") and immediately after transmembrane region 3 (IDRYLAIVHAVF; "DRY-box"). These two 12-amino acid segments are invariant among CCR1 through CCR4; CCR5 shares 15 22 of these 24 residues. These regions are sometimes conserved among other mammalian GPR and have been used for degenerate primer PCR to clone new CKRs. CRAM is divergent in these regions (9 out of 12 amino acids in TM2; 4 out of 12 in the DRY-box), which may explain why such approaches 20 have failed to identify CRAM. The DRY-box is in one of the three intracellular loops, and is thought to play a role in binding to heterotrimeric G proteins (Damaj, et al. (1996) FASEB J 10:1426). Because of the divergence of CRAM from the other  $\beta$ -CKRs in these regions, it may interact with a 25 different subset of G protein subunits, possibly transducing a signal different from that induced via other  $\beta$ -CKRs.

While human CKR genes have been localized to several different chromosomes, the  $\beta$ -CKR genes CCR1, CCR2, CCR3, 30 and CCR5 all cluster in a 350 kb region at chromosome 3p21.3 (Samson, et al. (1996) Genomics 36:522). CCR4 and the orphan receptors TER1 and GPR5 are also located in this 3p21 region Napolitano, et al. (1996) J Immunol 157:2759; Samson, et al. (1996) Genomics 36:522; Heiber, et al. 35 (1995) DNA Cell Biol 14:25) We determined the chromosomal location of CRAM. The Stanford G3 panel of radiation hybrids was used as templates for PCR reactions with CRAM-

specific primers. Among the 83 different hybrids, 11 contained the CRAM gene as assessed by PCR. CRAM co-localized with STS D3S3888, which is located at chromosome 3p21.3. Confirmation of this result was obtained from the recently completed sequence of the 143 kb BAC clone 110p12 from the 3p21 region (GenBank accession U95626); this BAC contains the loci CCR2, CCR5, and CRAM.

A related but different CRAM cDNA was also isolated from an activated monocyte library by random sequencing. Comparing the two forms of CRAM to the genomic sequence revealed the existence of two short exons (corresponding to 95750-96064 bp and 96186-96256 bp on BAC 110p12), followed by a large third exon (96630-98093 bp) that contains almost the entire ORF for CRAM. These two CRAM cDNAs consist of either exon 2 and exon 3 (1536 bp), or exon 1 and exon 3 (1780 bp). Exon 2 contributes 12 amino acids in frame with exon 3 to form the entire 356 residue polypeptide (CRAM-A). As exon 1 has no methionine in frame with the ORF in exon 3, the translated protein from this splice variant would start with Met-13, resulting in an N-terminally truncated protein of 344 amino acids (CRAM-B).

#### H. Expression of CRAM mRNA

RNA blot analysis showed expression largely restricted to lymphoid tissues. Prominent expression of CRAM mRNA was observed in spleen, lymph node, thymus, bone marrow, and fetal liver. Very little expression was detected in brain, liver, muscle, kidney, pancreas, or PBL, with moderate signals in heart, placenta, lung, and appendix. This pattern of expression was similar to that of the CKR-like gene *TER1* (Napolitano, et al. (1996) J Immunol 157:2759), but quite different from the related orphan receptor genes *V28* and *blr1* (Forster, et al. (1996) Cell 87:1037; and Raport, et al. (1995) Gene 163:295).

Data from various hematopoietic cell types showed no evidence for CRAM expression in resting or activated lymphocytes, or in splenocytes. CRAM mRNA was also not detected in resting monocytic cell lines, but was strongly

expressed in primary monocytes and THP-1 cells upon activation with IFN $\gamma$  and LPS. Both CRAM-A and CRAM-B mRNAs were induced, as detected by RT-PCR using exon 1- and exon 2-specific primers. In contrast to several other monocyte activation-induced genes, such as monokines (TNF $\alpha$ , IL-1, IL-6) and some cell-surface antigens (Ho and Moore. (1994) Therapeutic Immunology 1:173). CRAM mRNA expression was not detectably inhibited by IL-10. Thus, CRAM expression in monocytes may be regulated via a different mechanism compared to that of several other activation-induced genes.

While most CKR genes lack introns, the genes for human CCR2 and mouse CXCR4 (fusin) both contain at least two exons and both have two alternatively spliced forms. CCR2A and CCR2B differ in the C-terminus (Charo, et al. (1994) Proc. Natl. Acad. Sci. U.S.A. 91:2752), whereas CXCR4 has two forms that differ by two amino acids at the N-terminus (Heesen, et al. (1997) J. Immunol. 158:3561). The two forms of CCR2 have identical ligand specificities, but differ with respect to which G $\alpha$  subunits they can couple (Kuang, et al. (1996) J. Biol. Chem. 271:3975); the two forms of CXCR4 can both serve as functional CKRs for SDF-1 $\alpha$  (Heesen, et al. (1997) J. Immunol. 158:3561), although their ligand specificities and interactions with HIV have not been fully characterized. The N-terminal sequence of CKRs, along with portions of the extracellular loops, is known to play a key role in ligand binding and possibly receptor activation (Ahuja, et al. (1996) J. Biol. Chem. 271:225; Lu, et al. (1995) J. Biol. Chem. 270:26239; Horuk (1994) Immunol. Today 15:169; Wells, et al. (1996) J Leukoc Biol 59:53; and Hebert, et al. (1993) J. Biol. Chem. 268:18549). This region of CKR is also important for HIV fusion (Rucker, et al. (1996) Cell 87:437.), which is antagonized by chemokine ligands (Paxton, et al. (1996) Nature Med. 2:412; and Cocchi, et al. (1995) Science 270:1811). Thus it is possible that CRAM-A and CRAM-B may exhibit different but likely overlapping ligand specificities. Regulated expression of alternative forms

of a single CKR, combined with possible modulation of specificity of ligand-receptor interaction by chemokine-proteoglycan interaction (Graham, et al. (1996) The EMBO J. 15:6506; and Witt and Lander (1994) Curr. Biol. 4:394),

5 might control the spectrum of chemokines to which a particular cell could respond. In addition, these observations may provide one possible explanation of non-reciprocal desensitization phenomena observed with, for example, the chemokines RANTES, MIP-1 $\alpha$ , and MCAF (Wang, et  
10 al. (1993) J Exp Med 177:699).

The similarity of CRAM to the other  $\beta$ -CKRs, its chromosomal localization in the  $\beta$ -CKR gene cluster, and induction of its expression in monocytes upon activation all argue that CRAM may play an important role in  
15 regulation of immune function.

#### VIII. Screening for receptor/ligand

Labeled reagent is useful for screening of an  
20 expression library made from a cell line which expresses a chemokine or receptor, as appropriate. Standard staining techniques are used to detect or sort intracellular or surface expressed ligand, or surface expressing transformed cells are screened by panning. Screening of intracellular  
25 expression is performed by various staining or immunofluorescence procedures. See also, e.g., McMahan, et al. (1991) EMBO J. 10:2821-2832.

For example, on day 0, precoat 2-chamber permanox slides with 1 ml per chamber of fibronectin, 10 ng/ml in  
30 PBS, for 30 min at room temperature. Rinse once with PBS. Then plate COS cells at 2-3 x 10<sup>5</sup> cells per chamber in 1.5 ml of growth media. Incubate overnight at 37° C.

On day 1 for each sample, prepare 0.5 ml of a solution of 66  $\mu$ g/ml DEAE-dextran, 66  $\mu$ M chloroquine, and 4  $\mu$ g DNA  
35 in serum free DME. For each set, a positive control is prepared, e.g., of huIL-10-FLAG cDNA at 1 and 1/200 dilution, and a negative mock. Rinse cells with serum free

DME. Add the DNA solution and incubate 5 hr at 37° C. Remove the medium and add 0.5 ml 10% DMSO in DME for 2.5 min. Remove and wash once with DME. Add 1.5 ml growth medium and incubate overnight.

5 On day 2, change the medium. On days 3 or 4, the cells are fixed and stained. Rinse the cells twice with Hank's Buffered Saline Solution (HBSS) and fix in 4% paraformaldehyde (PFA)/glucose for 5 min. Wash 3X with HBSS. The slides may be stored at -80° C after all liquid  
10 is removed. For each chamber, 0.5 ml incubations are performed as follows. Add HBSS/saponin(0.1%) with 32 µl/ml of 1M NaN<sub>3</sub> for 20 min. Cells are then washed with HBSS/saponin 1X. Add antibody complex to cells and incubate for 30 min. Wash cells twice with HBSS/saponin.  
15 Add second antibody, e.g., Vector anti-mouse antibody, at 1/200 dilution, and incubate for 30 min. Prepare ELISA solution, e.g., Vector Elite ABC horseradish peroxidase solution, and preincubate for 30 min. Use, e.g., 1 drop of solution A (avidin) and 1 drop solution B (biotin) per 2.5  
20 ml HBSS/saponin. Wash cells twice with HBSS/saponin. Add ABC HRP solution and incubate for 30 min. Wash cells twice with HBSS, second wash for 2 min, which closes cells. Then add Vector diaminobenzoic acid (DAB) for 5 to 10 min. Use 2 drops of buffer plus 4 drops DAB plus 2 drops of H<sub>2</sub>O<sub>2</sub> per  
25 5 ml of glass distilled water. Carefully remove chamber and rinse slide in water. Air dry for a few minutes, then add 1 drop of Crystal Mount and a cover slip. Bake for 5 min at 85-90° C.

Alternatively, the binding compositions are used to  
30 affinity purify or sort out cells expressing the ligand or receptor. See, e.g., Sambrook, et al. or Ausubel et al.

All references cited herein are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and  
35 individually indicated to be incorporated by reference.

Many modification an variations of this invention can be made without departing from its spirit and scope, as

will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only, and the invention is to be limited only by the terms of the appended claims, along with the full scope of the  
5 equivalents to which such claims are entitled.